

Positivity Problems for Low-Order Linear Recurrence Sequences*

Joël Ouaknine[†]

James Worrell[‡]

Abstract

We consider two decision problems for linear recurrence sequences (LRS) over the integers, namely the *Positivity Problem* (are all terms of a given LRS positive?) and the *Ultimate Positivity Problem* (are all but finitely many terms of a given LRS positive?). We show decidability of both problems for LRS of order 5 or less, with complexity in the Counting Hierarchy for Positivity, and in polynomial time for Ultimate Positivity. Moreover, we show by way of hardness that extending the decidability of either problem to LRS of order 6 would entail major breakthroughs in analytic number theory, more precisely in the field of Diophantine approximation of transcendental numbers.

1 Introduction

A (real) **linear recurrence sequence (LRS)** is an infinite sequence $\mathbf{u} = \langle u_0, u_1, u_2, \dots \rangle$ of real numbers having the following property: there exist constants a_1, a_2, \dots, a_k (with $a_k \neq 0$) such that, for all $n \geq 0$,

$$(1.1) \quad u_{n+k} = a_1 u_{n+k-1} + a_2 u_{n+k-2} + \dots + a_k u_n.$$

If the initial values u_0, \dots, u_{k-1} of the sequence are provided, the recurrence relation defines the rest of the sequence uniquely. Such a sequence is said to have **order k** .¹

The best-known example of an LRS was given by Leonardo of Pisa in the 12th century: the Fibonacci sequence $\langle 0, 1, 1, 2, 3, 5, 8, 13, \dots \rangle$, which satisfies the recurrence relation $u_{n+2} = u_{n+1} + u_n$. Leonardo of Pisa introduced this sequence as a means to model the growth of an idealised population of rabbits. Not only has the Fibonacci sequence been extensively studied since, but LRS now form a vast subject in their own right, with numerous applications in mathematics and other sciences. A deep and extensive treatise on the

mathematical aspects of recurrence sequences is the recent monograph of Everest *et al.* [26].

In this paper, we focus on two key *decision problems* for LRS over the integers (or equivalently, for our purposes, the rationals):

- The **Positivity Problem**: given an LRS \mathbf{u} , are all terms of \mathbf{u} positive?
- The **Ultimate Positivity Problem**: given an LRS \mathbf{u} , are all but finitely many terms of \mathbf{u} positive?²

These problems (and assorted variants) have applications in a wide array of scientific areas, such as theoretical biology (analysis of L-systems, population dynamics) [36], economics (stability of supply-and-demand equilibria in cyclical markets, multiplier-accelerator models) [6], software verification (termination of linear programs) [45, 56, 17, 23, 16, 11], probabilistic model checking (reachability and approximation in Markov chains, stochastic logics) [7, 1], quantum computing (threshold problems for quantum automata) [13, 25], discrete linear dynamical systems (reachability and invariance problems) [32, 54, 10, 20], as well as combinatorics, formal languages, statistical physics, generating functions, etc. For example, as discussed in [38], terms of an LRS usually have combinatorial significance only if they are positive. Likewise, an LRS modelling population size is biologically meaningful only if it is uniformly positive.

Both Positivity and Ultimate Positivity bear some relationship to the well-known *Skolem Problem*: does a given LRS have a zero? The decidability of the Skolem Problem is generally considered to have been open since the 1930s (notwithstanding the fact that algorithmic decision issues had not at the time acquired the importance that they have today—see [30] for a discussion on this subject; see also [53] and [37], in which this state of affairs—the enduring openness of decidability for the Skolem Problem—is described as “faintly outrageous”

*This research was partially supported by EPSRC. We are also grateful to Matt Daws for considerable assistance in the initial stages of this work.

[†]Department of Computer Science, Oxford University, UK.

[‡]Department of Computer Science, Oxford University, UK.

¹Some authors define the order of an LRS as the *least* k such that the LRS obeys such a recurrence relation. The definition we have chosen allows for a simpler presentation of our results and is algorithmically more convenient.

²Note that both problems come in two natural flavours, according to whether strict or non-strict positivity is required. This paper focusses on the non-strict version, but alternatives and extensions (including strictness) are discussed in Section 6.

by Tao and a “mathematical embarrassment” by Lipton). A breakthrough occurred in the mid-1980s, when Mignotte *et al.* [41] and Vereshchagin [57] independently showed decidability for real algebraic LRS of order 4 or less. These deep results make essential use of Baker’s theorem on linear forms in logarithms (which earned Baker the Fields medal in 1970), as well as a p -adic analogue of Baker’s theorem due to van der Poorten. Unfortunately, little progress on that front has since been recorded.³ The Skolem Problem is known to be NP-hard if the order is unrestricted [14].

It is considered folklore that the decidability of Positivity would entail that of the Skolem Problem (see Section 2), noting however that the reduction increases the order of LRS quadratically. Nevertheless, the earliest explicit references in the literature to the Positivity and Ultimate Positivity Problems that we have found are from the 1970s (see, e.g., [51, 49, 12]). In [51], the Skolem and Positivity Problems are described as “very difficult”, whereas in [48], the authors assert that the Skolem, Positivity, and Ultimate Positivity Problems are “generally conjectured [to be] decidable”. Positivity and/or Ultimate Positivity are again stated as open in [29, 8, 35, 38, 54, 52], among others.

Unsurprisingly, progress on the Positivity and Ultimate Positivity Problems has been fairly slow. In the early 1980s, Burke and Webb showed that Ultimate Positivity is decidable for LRS of order 2 [18], and nine years later Nagasaka and Shiue [42] showed the same for LRS of order 3 that have repeated characteristic roots. Much more recently, Halava *et al.* showed that Positivity is decidable for integer LRS of order 2 [29], and three years later Laohakosol and Tangsupphathawat proved that both Positivity and Ultimate Positivity are decidable for integer LRS of order 3 [35]. In 2012, an article claiming to show decidability of Positivity for LRS of order 4 was published [52], with the authors noting being unable to tackle the case of order 5. Unfortunately, as acknowledged subsequently by the authors themselves [34], that paper contains a major error (the purported proof of Claim 2 on p.141, aimed at handling the most difficult critical case at order 4, is wrong, and appears not to be fixable without making use of sophisticated tools from analytic number theory as is done in the present paper).

To the best of our knowledge, no complexity bounds

³A proof of decidability of the Skolem Problem for LRS of order 5 was announced in [30]. However, as pointed out in [43], the proof seems to have a serious gap.

It is worth remarking, on the other hand, that whether an integer LRS has *infinitely* many zeros is known to be decidable at all orders [12].

currently exist in the literature concerning either the Positivity or Ultimate Positivity Problems, other than coNP-hardness for LRS of unbounded orders which is inherited from the reduction from the Skolem Problem (cf. Section 2).

Our main results are as follows:⁴

- The Positivity Problem is decidable for integer LRS of order 5 or less, with complexity in $\text{coNP}^{\text{PP}^{\text{PP}^{\text{PP}}}}$, i.e., within the fourth level of the Counting Hierarchy.
- The Ultimate Positivity Problem is decidable for integer LRS of order 5 or less in polynomial time.
- The decidability of either Positivity or Ultimate Positivity for integer LRS of order 6 would entail major breakthroughs in analytic number theory (certain open problems in Diophantine approximation of transcendental numbers long believed to be hard would become solvable)—see Section 5 for precise statements.

These results, which—absent major advances in number theory—can essentially be viewed as completing the picture on Positivity problems for linear recurrence sequences, substantially improve the state of the art over the last three decades’ worth of research on the subject. Most prior work on Positivity problems that we are aware of has been confined to the use of linear algebra and elementary algebraic number theoretic techniques. By contrast, we are deploying in this paper an eclectic arsenal of deep and sophisticated mathematical tools from analytic and algebraic number theory, Diophantine geometry and approximation, and real algebraic geometry, notably Baker’s theorem on linear forms in logarithms, Masser’s results on multiplicative relationships among algebraic numbers, Kronecker’s theorem on simultaneous Diophantine approximation, and Renegar’s work on the fine-grained complexity of the first-order theory of the reals. These results are summarised in Section 3. We then present a high-level overview of our proof strategy—split in two parts—in the first half of Section 4. Various extensions and generalisations of our results, along with avenues for future work, are discussed in Section 6.

⁴The complexities are given as a function of the bit length of standard representations of integer LRS of order k ; for an LRS as defined by Equation (1.1), this representation consists of the $2k$ -tuple $(a_1, \dots, a_k, u_0, \dots, u_{k-1})$ of integers.

Note also that the Counting-Hierarchy complexity class does not require parenthesising since $\text{co}(\text{NP}^{\text{PP}^{\text{PP}^{\text{PP}}}}) = (\text{coNP})^{\text{PP}^{\text{PP}^{\text{PP}}}}$.

2 Linear Recurrence Sequences

We recall some fundamental properties of linear recurrence sequences. Results are stated without proof, and we refer the reader to [26, 30] for details.

Let $\mathbf{u} = \langle u_n \rangle_{n=0}^\infty$ be an LRS of order k over the reals satisfying the recurrence relation

$$u_{n+k} = a_1 u_{n+k-1} + \dots + a_k u_n,$$

where without loss of generality we may assume that $a_k \neq 0$. We denote by $\|\mathbf{u}\|$ the bit length of its representation as a $2k$ -tuple of integers, as discussed in the previous section. The **characteristic polynomial** of \mathbf{u} is

$$p(x) = x^k - a_1 x^{k-1} - \dots - a_{k-1} x - a_k.$$

The **characteristic roots** of \mathbf{u} are the roots of this polynomial, and the **dominant roots** are the roots of maximum modulus.

The characteristic roots divide naturally into those that are real and those that are not. As we exclusively deal with LRS over the reals, the characteristic polynomial has real coefficients and non-real roots therefore always arise in conjugate pairs. Thus we may write $\{\rho_1, \dots, \rho_\ell, \gamma_1, \overline{\gamma_1}, \dots, \gamma_m, \overline{\gamma_m}\}$ to represent the set of characteristic roots of \mathbf{u} , where each $\rho_i \in \mathbb{R}$ and each $\gamma_j \in \mathbb{C} \setminus \mathbb{R}$. There are now univariate polynomials A_1, \dots, A_ℓ and C_1, \dots, C_m such that, for all $n \geq 0$,

$$u_n = \sum_{i=1}^{\ell} A_i(n) \rho_i^n + \sum_{j=1}^m (C_j(n) \gamma_j^n + \overline{C_j(n)} \overline{\gamma_j}^n).$$

This expression is referred to as the **exponential polynomial** solution of \mathbf{u} . For integer LRS, the polynomials A_i have real algebraic coefficients and the polynomials C_j have complex algebraic coefficients. The degree of each of these polynomials is at most one less than the multiplicity of the corresponding characteristic root; thus in particular, these polynomials are identically constant when \mathbf{u} has no repeated characteristic roots. For fixed k , all coefficients appearing in these polynomials can be computed in time polynomial in $\|\mathbf{u}\|$, since they can be obtained by solving a system of linear equations involving the first k terms of \mathbf{u} . See Section 3 for further details on algebraic-number manipulations.

An LRS is said to be **non-degenerate** if it does not have two distinct characteristic roots whose quotient is a root of unity. As pointed out in [26], the study of arbitrary LRS can effectively be reduced to that of non-degenerate LRS, by partitioning the original LRS into finitely many subsequences, each of which is non-degenerate. In general, such a reduction will require exponential time. However, when restricting ourselves

to LRS of bounded order (in our case, of order at most 5), the reduction can be carried out in polynomial time. In particular, any LRS of order 5 or less can be partitioned in polynomial time into at most 2520 non-degenerate LRS of the same order or less.⁵ In the rest of this paper, we shall therefore assume that all LRS we are given are non-degenerate.

Any LRS \mathbf{u} of order k can alternately be given in matrix form, in the sense that there is a square matrix M of dimension $k \times k$, together with k -dimensional column vectors \vec{v} and \vec{w} , such that, for all $n \geq 0$, $u_n = \vec{v}^T M^n \vec{w}$. It suffices to take M to be the transpose of the companion matrix of the characteristic polynomial of \mathbf{u} , let \vec{v} be the vector (u_{k-1}, \dots, u_0) of initial terms of \mathbf{u} in reverse order, and take \vec{w} to be the vector whose first $k-1$ entries are 0 and whose k th entry is 1. It is worth noting that the characteristic roots of \mathbf{u} correspond precisely to the eigenvalues of M . This translation is instrumental in Section 4 to place the Positivity Problem for LRS of order at most 5 within the Counting Hierarchy.

Conversely, given any square matrix M of dimension $k \times k$, and any k -dimensional vectors \vec{v} and \vec{w} , let $u_n = \vec{v}^T M^n \vec{w}$. Then $\langle \vec{v}^T M^n \vec{w} \rangle_{n=0}^\infty$ is an LRS of order at most k whose characteristic polynomial is the same as that of M , as can be seen by applying the Cayley-Hamilton Theorem.⁶

Let $\langle u_n \rangle_{n=0}^\infty$ and $\langle v_n \rangle_{n=0}^\infty$ be LRS of order k and l respectively. Their pointwise product $\langle u_n v_n \rangle_{n=0}^\infty$ and sum $\langle u_n + v_n \rangle_{n=0}^\infty$ are also LRS of order at most kl and $k+l$ respectively. In the special case of pointwise squaring, the order of the LRS $\langle u_n^2 \rangle_{n=0}^\infty$ is at most $k(k+1)/2$.

We can use the above to reduce (the complement of) the Skolem Problem to Positivity: given an integer LRS $\mathbf{u} = \langle u_n \rangle_{n=0}^\infty$, we see that $u_n \neq 0$ iff $u_n^2 - 1 \geq 0$. Since this reduction is polynomial in $\|\mathbf{u}\|$, the NP-hardness for the Skolem Problem presented in [14] immediately translates as coNP-hardness for Positivity, as pointed out in [9]. In fact, since the LRS used in [14] are all periodic, we also obtain a coNP-hardness for Ultimate Positivity. At the time of writing, no other complexity bounds for these problems are known.

3 Mathematical Tools

In this section we introduce the key technical tools used in this paper.

⁵We obtained this value using a bespoke enumeration procedure for order 5. A bound of $e^{2\sqrt{6.5 \log 5}} \leq 1,085,134$ can be obtained from Corollary 3.3 of [58].

⁶In fact, if none of the eigenvalues of M are zero, it is easy to see that the full sequence $\langle \vec{v}^T M^n \vec{w} \rangle_{n=0}^\infty$ is an LRS (of order at most k).

For $p \in \mathbb{Z}[x_1, \dots, x_m]$ a polynomial with integer coefficients, let us denote by $\|p\|$ the bit length of its representation as a list of coefficients encoded in binary. Note that the degree of p is at most $\|p\|$, and the height of p —i.e., the maximum of the absolute values of its coefficients—is at most $2^{\|p\|}$.

We begin by summarising some basic facts about algebraic numbers and their (efficient) manipulation. The main references include [21, 5, 46].

A complex number α is **algebraic** if it is a root of a single-variable polynomial with integer coefficients. The **defining polynomial** of α , denoted p_α , is the unique polynomial of least degree, and whose coefficients do not have common factors, which vanishes at α . The **degree** and **height** of α are respectively those of p_α .

A standard representation⁷ for algebraic numbers is to encode α as a tuple comprising its defining polynomial together with rational approximations of its real and imaginary parts of sufficient precision to distinguish α from the other roots of p_α . More precisely, α can be represented by $(p_\alpha, a, b, r) \in \mathbb{Z}[x] \times \mathbb{Q}^3$ provided that α is the unique root of p_α inside the circle in \mathbb{C} of radius r centred at $a + bi$. A separation bound due to Mignotte [40] asserts that for roots $\alpha \neq \beta$ of a polynomial $p \in \mathbb{Z}[x]$, we have

$$(3.2) \quad |\alpha - \beta| > \frac{\sqrt{6}}{d^{(d+1)/2} H^{d-1}},$$

where d and H are respectively the degree and height of p . Thus if r is required to be less than a quarter of the root-separation bound, the representation is well-defined and allows for equality checking. Given a polynomial $p \in \mathbb{Z}[x]$, it is well-known how to compute standard representations of each of its roots in time polynomial in $\|p\|$ [44, 21, 5]. Thus given α an algebraic number for which we have (or wish to compute) a standard representation, we write $\|\alpha\|$ to denote the bit length of this representation. From now on, when referring to computations on algebraic numbers, we always implicitly refer to their standard representations.

Note that Equation (3.2) can be used more generally to separate arbitrary algebraic numbers: indeed, two algebraic numbers α and β are always roots of the polynomial $p_\alpha p_\beta$ of degree at most the sum of the degrees of α and β , and of height at most the product of the heights of α and β .

Given algebraic numbers α and β , one can compute $\alpha + \beta$, $\alpha\beta$, $1/\alpha$ (for non-zero α), $\bar{\alpha}$, and $|\alpha|$, all of which are algebraic, in time polynomial in $\|\alpha\| + \|\beta\|$. Likewise, it is straightforward to check whether $\alpha = \beta$. Moreover, if $\alpha \in \mathbb{R}$, deciding whether $\alpha > 0$ can be done

⁷Note that this representation is not unique.

in time polynomial in $\|\alpha\|$. Efficient algorithms for all these tasks can be found in [21, 5].

Remarkably, integer multiplicative relationships among a fixed number of algebraic numbers can be elicited systematically in polynomial time:

THEOREM 3.1. *Let m be fixed, and let $\lambda_1, \dots, \lambda_m$ be complex algebraic numbers of modulus 1. Consider the free abelian group L under addition given by*

$$L = \{(v_1, \dots, v_m) \in \mathbb{Z}^m : \lambda_1^{v_1} \dots \lambda_m^{v_m} = 1\}.$$

L has a basis $\{\vec{\ell}_1, \dots, \vec{\ell}_p\} \subseteq \mathbb{Z}^m$ (with $p \leq m$), where the entries of each of the $\vec{\ell}_j$ are all polynomially bounded in $\|\lambda_1\| + \dots + \|\lambda_m\|$. Moreover, such a basis can be computed in time polynomial in $\|\lambda_1\| + \dots + \|\lambda_m\|$.

Note in the above that the bound is on the *magnitude* of the vectors $\vec{\ell}_j$ (rather than the bit length of their representation), which follows from a deep result of Masser [39]. For a proof of Theorem 3.1, see also [27, 19].

We now turn to the first-order theory of the reals. Let $\vec{x} = x_1, \dots, x_m$ be a list of m real-valued variables, and let $\sigma(\vec{x})$ be a Boolean combination of atomic predicates of the form $g(\vec{x}) \sim 0$, where each $g(\vec{x}) \in \mathbb{Z}[\vec{x}]$ is a polynomial with integer coefficients over these variables, and \sim is either $>$ or $=$. A **sentence of the first-order theory of the reals** is of the form

$$(3.3) \quad Q_1 x_1 \dots Q_m x_m \sigma(\vec{x}),$$

where each Q_i is one of the quantifiers \exists or \forall . Let us denote the above formula by τ , and write $\|\tau\|$ to denote the bit length of its syntactic representation.

Tarski famously showed that the first-order theory of the reals is decidable [55]. His procedure, however, has non-elementary complexity. Many substantial improvements followed over the years, starting with Collins's technique of cylindrical algebraic decomposition [22], and culminating with the fine-grained analysis of Renegar [46]. In this paper, we focus exclusively on the situation in which the number of variables is uniformly bounded.

THEOREM 3.2. (RENEGAR) *Let $M \in \mathbb{N}$ be fixed. Let τ be of the form (3.3) above. Assume that the number of variables in τ is bounded by M (i.e., $m \leq M$). Then the truth value of τ can be determined in time polynomial in $\|\tau\|$.*

Theorem 3.2 follows immediately from [46, Thm. 1.1].

Our next result is a special case of Kronecker's famous theorem on simultaneous Diophantine approximation, a statement and proof of which can be found in [15, Chap. 7, Sec. 1.3, Prop. 7].

For $x \in \mathbb{R}$, write $[x]_{2\pi}$ to denote the distance from x to the closest integer multiple of 2π : $[x]_{2\pi} = \min\{|x - 2\pi j| : j \in \mathbb{Z}\}$.

THEOREM 3.3. (KRONECKER) *Let $t_1, \dots, t_m, x_1, \dots, x_m \in [0, 2\pi)$. The following are equivalent:*

1. *For any $\varepsilon > 0$, there exists $n \in \mathbb{Z}$ such that, for $1 \leq j \leq m$, we have $[nt_j - x_j]_{2\pi} \leq \varepsilon$.*
2. *For every tuple (v_1, \dots, v_m) of integers such that $[v_1 t_1 + \dots + v_m t_m]_{2\pi} = 0$, we have $[v_1 x_1 + \dots + v_m x_m]_{2\pi} = 0$.*

We can strengthen Theorem 3.3 by requiring that $n \in \mathbb{N}$ in the first assertion. Indeed, suppose that in a given instance, we find that $n < 0$. A straightforward pigeonhole argument shows that there exist arbitrarily large positive integers g such that $[gt_j]_{2\pi} \leq \varepsilon$ for $1 \leq j \leq m$. It follows that $[(g+n)t_j - x_j]_{2\pi} \leq 2\varepsilon$, which establishes the claim for sufficiently large g (noting that ε is arbitrary).

Let $\lambda_1, \dots, \lambda_m$ be complex algebraic numbers of modulus 1. For each $j \in \{1, \dots, m\}$, write $\lambda_j = e^{i\theta_j}$ for some $\theta_j \in [0, 2\pi)$. Let

$$L = \{(v_1, \dots, v_m) \in \mathbb{Z}^m : \lambda_1^{v_1} \dots \lambda_m^{v_m} = 1\} \\ = \{(v_1, \dots, v_m) \in \mathbb{Z}^m : [v_1 \theta_1 + \dots + v_m \theta_m]_{2\pi} = 0\}.$$

Recall from Theorem 3.1 that L is a free abelian group under addition with basis $\{\vec{\ell}_1, \dots, \vec{\ell}_p\} \subseteq \mathbb{Z}^m$, where $p \leq m$.

For each $j \in \{1, \dots, p\}$, let $\vec{\ell}_j = (\ell_{j,1}, \dots, \ell_{j,m})$. Write

$$R = \{\vec{x} = (x_1, \dots, x_m) \in [0, 2\pi)^m : \\ [\vec{\ell}_j \cdot \vec{x}]_{2\pi} = 0 \text{ for } 1 \leq j \leq p\}.$$

By Theorem 3.3, for an arbitrary tuple $(x_1, \dots, x_m) \in [0, 2\pi)^m$, it is the case that, for all $\varepsilon > 0$, there exists $n \in \mathbb{N}$ such that, for $j \in \{1, \dots, m\}$, $[n\theta_j - x_j]_{2\pi} \leq \varepsilon$ iff $(x_1, \dots, x_m) \in R$.

Now observe that $(x_1, \dots, x_m) \in R$ iff $(e^{ix_1}, \dots, e^{ix_m}) \in T$, where

$$T = \{(z_1, \dots, z_m) \in \mathbb{C}^m : |z_1| = \dots = |z_m| = 1 \text{ and,} \\ \text{for each } j \in \{1, \dots, p\}, z_1^{\ell_{j,1}} \dots z_m^{\ell_{j,m}} = 1\}.$$

Since $e^{in\theta_j} = \lambda_j^n$, we immediately have the following:

COROLLARY 3.1. *Let $\lambda_1, \dots, \lambda_m$ and T be as above. Then $\{(\lambda_1^n, \dots, \lambda_m^n) : n \in \mathbb{N}\}$ is a dense subset of T .*

Finally, we give a version of Baker's deep theorem on linear forms in logarithms. The particular statement we have chosen is a sharp formulation due to Baker and Wüstholz [4].

In what follows, \log refers to the principal value of the complex logarithm function given by $\log z = \log |z| + i \arg z$, where $-\pi < \arg z \leq \pi$.

THEOREM 3.4. (BAKER AND WÜSTHOLZ) *Let $\alpha_1, \dots, \alpha_m \in \mathbb{C}$ be algebraic numbers different from 0 or 1, and let $b_1, \dots, b_m \in \mathbb{Z}$ be integers. Write*

$$\Lambda = b_1 \log \alpha_1 + \dots + b_m \log \alpha_m.$$

Let $A_1, \dots, A_m, B \geq e$ be real numbers such that, for each $j \in \{1, \dots, m\}$, A_j is an upper bound for the height of α_j , and B is an upper bound for $|b_j|$. Let d be the degree of the extension field $\mathbb{Q}(\alpha_1, \dots, \alpha_m)$ over \mathbb{Q} .

If $\Lambda \neq 0$, then

$$\log |\Lambda| > -(16md)^{2(m+2)} \log A_1 \dots \log A_m \log B.$$

Finally, we record the following fact, whose straightforward proof is left to the reader.

PROPOSITION 3.1. *Let $a \geq 2$ and $\varepsilon \in (0, 1)$ be real numbers. Let $B \in \mathbb{Z}[x]$ have degree at most a^{D_1} and height at most $2^{a^{D_2}}$, and assume that $1/\varepsilon \leq 2^{a^{D_3}}$, for some $D_1, D_2, D_3 \in \mathbb{N}$. Then there is $D_4 \in \mathbb{N}$ depending only on D_1, D_2, D_3 such that, for all $n \geq 2^{a^{D_4}}$, $\frac{1}{B(n)} > (1 - \varepsilon)^n$.*

4 Decidability and Complexity

Let $\mathbf{u} = \langle u_n \rangle_{n=0}^\infty$ be an integer LRS of order k . As discussed in the Introduction, we assume that u is presented as a $2k$ -tuple of integers $(a_1, \dots, a_k, u_0, \dots, u_{k-1}) \in \mathbb{Z}^{2k}$, such that for all $n \geq 0$,

$$(4.4) \quad u_{n+k} = a_1 u_{n+k-1} + \dots + a_k u_n.$$

The **Positivity Problem** asks, given such an LRS \mathbf{u} , whether for all $n \geq 0$, it is the case that $u_n \geq 0$. When this holds, we say that \mathbf{u} is **positive**.

The **Ultimate Positivity Problem** asks, given such an LRS \mathbf{u} , whether there exists $N \geq 0$ such that, for all $n \geq N$, it is the case that $u_n \geq 0$. When this holds, we say that \mathbf{u} is **ultimately positive**.

In this section, we establish the following main results:

THEOREM 4.1. *The Positivity Problem for integer LRS of order 5 or less is decidable in $\text{coNP}^{\text{PP}^{\text{PP}^{\text{PP}}}}$.*

THEOREM 4.2. *The Ultimate Positivity Problem for integer LRS of order 5 or less is decidable in polynomial time.*

Note that the above results immediately carry over to rational LRS. To see this, consider a rational LRS \mathbf{u} obeying the recurrence relation (4.4). Let ℓ be the least common multiple of the denominators of the rational numbers $a_1, \dots, a_k, u_0, \dots, u_{k-1}$, and define an integer sequence $\mathbf{v} = \langle v_n \rangle_{n=0}^\infty$ by setting $v_n = \ell^{n+1} u_n$ for all $n \geq 0$. It is easily seen that \mathbf{v} is an integer LRS of the same order as \mathbf{u} , and that for all n , $v_n \geq 0$ iff $u_n \geq 0$.

Positivity—High-Level Synopsis. At a high level, the algorithm upon which Theorem 4.1 rests proceeds as follow. Given an LRS \mathbf{u} , we first decide whether or not \mathbf{u} is ultimately positive by studying its exponential polynomial solution—further details on this task are provided shortly. As we prove in this paper, whenever \mathbf{u} is an ultimately positive LRS of order 5 or less, there is an effective bound N of at most exponential magnitude such that all terms of \mathbf{u} beyond N are positive. Next, observe that \mathbf{u} cannot be positive unless it is ultimately positive. Now in order to assert that an ultimately positive LRS \mathbf{u} is *not* positive, we use a *guess-and-check* procedure: find $n \leq N$ such that $u_n < 0$. By writing $u_n = \vec{v}^T M^n \vec{w}$, for some square integer matrix M and vectors \vec{v} and \vec{w} (cf. Section 2), we can decide whether $u_n < 0$ in PosSLP⁸ via iterative squaring, which yields an NP^{PosSLP} procedure for non-Positivity. Thanks to the work of Allender *et al.* [2], which asserts that PosSLP \subseteq P^{PPPP}, we obtain the required coNP^{PPPP} algorithm for deciding Positivity.

The following is an old result concerning LRS; proofs can be found in [28, Thm. 7.1.1] and [8, Thm. 2]. It also follows easily and directly from either Pringsheim’s theorem or from [17, Lem. 4]. It plays an important role in our approach by enabling us to significantly cut down on the number of subcases that must be considered, avoiding the sort of quagmire alluded to in [42].

PROPOSITION 4.1. *Let $\langle u_n \rangle_{n=0}^\infty$ be an LRS with no real positive dominant characteristic root. Then there are infinitely many n such that $u_n < 0$ and infinitely many n such that $u_n > 0$.*

By Proposition 4.1, it suffices to restrict our attention to LRS whose dominant characteristic roots include one real positive value. Given an integer LRS \mathbf{u} , note that determining whether the latter holds is easily done in time polynomial in $\|\mathbf{u}\|$.

Thus let \mathbf{u} be a non-degenerate integer LRS of order k having a (possibly repeated) real positive dom-

inant characteristic root $\rho > 0$. Note that \mathbf{u} cannot have a real negative dominant characteristic root (which would be $-\rho$), since otherwise the quotient $-\rho/\rho = -1$ would be a root of unity, contradicting non-degeneracy. Let us therefore write the characteristic roots as $\{\rho, \gamma_1, \overline{\gamma_1}, \dots, \gamma_m, \overline{\gamma_m}\} \cup \{\gamma_{m+1}, \gamma_{m+2}, \dots, \gamma_\ell\}$, where we assume that the roots in the first set all have common modulus ρ , whereas the roots in the second set all have modulus strictly smaller than ρ . Note that for LRS of order at most 5, m can be at most 2.

Let $\lambda_i = \gamma_i/\rho$ for $1 \leq i \leq \ell$. We can then write

$$(4.5) \quad \frac{u_n}{\rho^n} = A(n) + \sum_{i=1}^m \left(C_i(n) \lambda_i^n + \overline{C_i(n)} \overline{\lambda_i}^n \right) + r(n),$$

for a suitable real polynomial A and complex polynomials C_1, \dots, C_m , where $r(n)$ is a term tending to zero exponentially fast.

Note that none of $\lambda_1, \dots, \lambda_m$, all of which have modulus 1, can be a root of unity, as each λ_i is a quotient of characteristic roots and \mathbf{u} is assumed to be non-degenerate.

For $i \in \{1, \dots, \ell\}$, observe also that as each λ_i is a quotient of two roots of the same polynomial of degree k , it has degree at most $k(k-1)$. In fact, it is easily seen that $\|\lambda_i\| = \|\mathbf{u}\|^{\mathcal{O}(1)}$.

As noted in Section 2, the degree of polynomials A and C_i in the exponential polynomial solution is at most one less than the multiplicity of the corresponding characteristic roots, and is therefore bounded above by $k-1$. Recall also that all coefficients appearing in these polynomials are algebraic and, for fixed k , can be computed and manipulated in time polynomial in $\|\mathbf{u}\|$. It easily follows that $\|A\| = \|\mathbf{u}\|^{\mathcal{O}(1)}$ and $\|C_i\| = \|\mathbf{u}\|^{\mathcal{O}(1)}$.

Finally, we place bounds on the rate of convergence of $r(n)$. We have

$$r(n) = C_{m+1}(n) \lambda_{m+1}^n + \dots + C_\ell(n) \lambda_\ell^n.$$

For fixed k , combining our estimates on the height and degree of each λ_i together with the root-separation bound given by Equation (3.2), we get $\left| \frac{1}{1-\lambda_i} \right| = 2\|\mathbf{u}\|^{\mathcal{O}(1)}$, for $m+1 \leq i \leq \ell$. Thanks also to the bounds on the height and degree of the polynomials C_i , it follows that we can find $\varepsilon \in (0, 1)$ and $N \in \mathbb{N}$ such that:

$$(4.6) \quad 1/\varepsilon = 2\|\mathbf{u}\|^{\mathcal{O}(1)}$$

$$(4.7) \quad N = 2\|\mathbf{u}\|^{\mathcal{O}(1)}$$

$$(4.8) \quad \text{For all } n > N, |r(n)| < (1 - \varepsilon)^n.$$

In addition, we can compute such ε and N in time polynomial in $\|\mathbf{u}\|$. Naturally, given k , we can also

⁸Recall that PosSLP is the problem of determining whether an arithmetic circuit, with addition, multiplication, and subtraction gates, evaluates to a positive integer.

assume that we have calculated explicitly once and for all the constants implicit in the various instances of the $\mathcal{O}(1)$ notation.

We now seek to answer Positivity and Ultimate Positivity for the LRS $\mathbf{u} = \langle u_n \rangle_{n=0}^\infty$ by studying the same for $\langle u_n / \rho^n \rangle_{n=0}^\infty$.

In what follows, we assume that \mathbf{u} is as given above; in particular, \mathbf{u} is a non-degenerate integer LRS having a (possibly repeated) real positive dominant characteristic root $\rho > 0$.

Ultimate Positivity—High-Level Synopsis. Before launching into technical details, let us provide a high-level overview of our proof strategy for deciding Ultimate Positivity. Consider first the special case of Equation (4.5) in which the polynomials $A(n)$ and $C_i(n), \overline{C}_i(n)$ are all identically constant. Let us rewrite this equation as

$$(4.9) \quad \frac{u_n}{\rho^n} = A + h(\lambda_1^n, \dots, \lambda_m^n) + r(n),$$

where $h : \mathbb{C}^m \rightarrow \mathbb{R}$ is a continuous function. In general, there will be integer multiplicative relationships among the $\lambda_1, \dots, \lambda_m$, for which we can compute a basis B thanks to Theorem 3.1. These multiplicative relationships define a torus $T \subseteq \mathbb{C}^m$ on which the joint iterates $(\lambda_1^n, \dots, \lambda_m^n)$ are dense, as per Kronecker's theorem (in the form of Corollary 3.1).

If $r(n)$ is identically 0, then both Positivity and Ultimate Positivity can be decided by determining the sign of the expression $A + \min h|_T$ (where $h|_T$ denotes the function h restricted to the torus T). For fixed order k , computing this sign can be carried out in polynomial time via the first-order theory of the reals, thanks to Theorem 3.2.

If $r(n)$ is not identically 0, then for LRS of order at most 5, we have that m is either 0 or 1, where the latter is the interesting case. The torus T is now simply the unit circle in the complex plane, and Equations (4.5) and (4.9) can be rewritten as

$$\frac{u_n}{\rho^n} = A + 2|c_1| \cos(n\theta_1 + \varphi_1) + r(n),$$

where $C_1(n) = c_1 = |c_1|e^{i\varphi_1}$ and $\theta_1 = \arg \lambda_1$. The critical case now arises when $A - 2|c_1| = 0$, which we can determine in polynomial time. Noting that the cosine function is minimised when its argument is an odd integer multiple of π , we can use Baker's theorem to bound the expression $n\theta_1 + \varphi_1$ away from odd integer multiples of π by an inverse polynomial in n . Using a Taylor approximation, we then argue that $\cos(n\theta_1 + \varphi_1)$ is itself eventually bounded away from -1 by a (different) inverse polynomial in n , and

since $r(n)$ decays to zero exponentially fast, we are able to conclude that u_n / ρ^n is ultimately positive, and can compute a bound N after which all terms u_n (for $n > N$) are positive.

Returning to Equation (4.5), note that if the $C_i(n), \overline{C}_i(n)$ are all identically constant but $A(n)$ is not, then the latter will eventually dominate and enable us to settle the ultimate positivity question; likewise, if $A(n)$ is identically constant but some $C_i(n), \overline{C}_i(n)$ are not, the latter eventually dominate and the situation can be dealt with straightforwardly.

This analysis allows us to handle LRS of order up to 5. At order 6, however, we encounter a critical situation in which $A(n)$ and $C_1(n), \overline{C}_1(n)$ are all linear polynomials, which then leads to the hardness results described in Section 5.

We now proceed with the proofs of Theorems 4.1 and 4.2, split into cases according to the number of distinct (albeit possibly repeated) dominant characteristic roots of \mathbf{u} . Since there is one real positive dominant root, no real negative dominant root, and since non-real roots always arise in pairs, the number of dominant roots must be odd. In any event, the total number of characteristic roots is bounded by the order of \mathbf{u} , which we assume to be at most 5.

4.1 One Dominant Root. In case of a single dominant root $\rho \in \mathbb{R}$, from Equation (4.5) we have that $u_n / \rho^n = A(n) + r(n)$. If $A(n)$ is identically 0, we simply turn our attention towards $r(n)$, which is an LRS whose exponential polynomial solution has one fewer term. Otherwise, it is clear that \mathbf{u} is ultimately positive iff either $A(n)$ is identically equal to some constant $a > 0$, or $\lim_{n \rightarrow \infty} A(n) = \infty$, all of which can be decided straightforwardly in time polynomial in $\|\mathbf{u}\|$.

Turning to positivity, assume therefore that $A(n)$ is either a strictly positive constant or tends to ∞ . Recall from our earlier discussion on the rate of convergence of $r(n)$ that we can compute in polynomial time numbers $\varepsilon \in (0, 1)$, with $1/\varepsilon = 2^{\|\mathbf{u}\|^{O(1)}}$, and $N = 2^{\|\mathbf{u}\|^{O(1)}}$, such that $|r(n)| < (1 - \varepsilon)^n$ for all $n > N$. We can similarly compute a bound $N' = 2^{\|\mathbf{u}\|^{O(1)}}$ such that $A(n) \geq (1 - \varepsilon)^n$ for all $n > N'$. Let $N'' = \max\{N, N'\}$. Then \mathbf{u} will fail to be positive iff there is some $n \leq N''$ such that $u_n < 0$. Since N'' is at most exponential in $\|\mathbf{u}\|$, we can decide positivity of \mathbf{u} in $\text{coNP}^{\text{PP}^{\text{PP}^{\text{PP}}}}$ via a PosSLP oracle as outlined earlier.

4.2 Three Dominant Roots. Next, we consider the case in which \mathbf{u} has exactly three dominant characteristic roots $\{\rho, \gamma_1, \overline{\gamma}_1\}$. Two subcases arise: (i) either the

complex roots γ_1 and $\overline{\gamma_1}$ are simple, or (ii) γ_1 and $\overline{\gamma_1}$ are repeated.

(i) In the first subcase, the multiplicity of ρ may range from 1 to 3. If ρ has multiplicity 3 then there can be no other characteristic roots, and

$$u_n/\rho^n = an^2 + bn + d + c_1\lambda_1^n + \overline{c_1}\overline{\lambda_1}^n,$$

where a, b, d are real algebraic constants, and c_1 is a complex algebraic constant which we assume is non-zero (otherwise the situation is trivial).

If $a < 0$, then clearly \mathbf{u} is neither positive nor ultimately positive. If $a > 0$ then \mathbf{u} is ultimately positive and, similarly to the case of a single dominant root, we can use our earlier estimates on the height and degree of a, b, d , and c_1 , together with the root-separation bound given by Equation (3.2), to conclude that there is $N = 2^{\|\mathbf{u}\|^{O(1)}}$ such that, for all $n > N$, we have $u_n \geq 0$. The positivity of \mathbf{u} can then be decided in $\text{coNP}^{\text{PPP}^{\text{PPP}}}$.

Next, if $a = 0$ then there is potentially an exponentially decaying term in the exponential polynomial solution of u_n/ρ^n ; this also covers the case in which the multiplicity of ρ is 1 or 2:

$$u_n/\rho^n = bn + d + c_1\lambda_1^n + \overline{c_1}\overline{\lambda_1}^n + r(n).$$

Here, similarly to the previous case, if $b < 0$ then \mathbf{u} is neither positive nor ultimately positive, whereas if $b > 0$ then \mathbf{u} is ultimately positive and, as before, we obtain an exponential upper bound on the index n of possible violations of positivity, as required.

Finally, suppose that $a = 0$ and $b = 0$. We may assume that $c_1 \neq 0$, otherwise we are left with the term $r(n)$ and can simply recast our analysis appropriately at lower order. Let $\theta_1 = \arg \lambda_1$ and $\varphi_1 = \arg c_1$. We have

$$\frac{u_n}{\rho^n} = d + 2|c_1| \cos(n\theta_1 + \varphi_1) + r(n).$$

Since λ_1 is not a root of unity, it is straightforward to see that the set $\{\cos(n\theta_1 + \varphi_1) : n \geq 0\}$ is dense in $[-1, 1]$. It immediately follows that if $d < 2|c_1|$ then \mathbf{u} is neither positive nor ultimately positive, whereas if $d > 2|c_1|$ then \mathbf{u} is ultimately positive with, as before, an exponential bound on the index of possible violations of positivity.

It remains to tackle the case in which $d = 2|c_1|$. Since λ_1 is not a root of unity, there is at most one value of n such that $n\theta_1 + \varphi_1$ is an odd integer multiple of π , corresponding to $\lambda_1^n = -|c_1|/c_1$. It then follows from Theorem 3.1 that this value (if it exists) is at most $M = \|\mathbf{u}\|^{O(1)}$.

By Equations (4.6)–(4.8), we can find $\varepsilon \in (0, 1)$ and $N = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for all $n > N$, we have $|r(n)| < (1 - \varepsilon)^n$, and moreover $1/\varepsilon = 2^{\|\mathbf{u}\|^{O(1)}}$.

Let $g(x) = \frac{x^2}{2!} - \frac{x^4}{4!}$. Using a Taylor approximation, we have the following:

$$(4.10) \quad \cos(x + \pi) \geq -1 + g(x) \quad \text{for } x \in (-\pi, \pi]$$

$$(4.11) \quad g(x) \leq g(y) \quad \text{for } |x| \leq |y| \leq 1$$

$$(4.12) \quad 11/24 = g(1) \leq g(x) \quad \text{for } 1 \leq |x| \leq \pi.$$

For $n \in \mathbb{N}$, write $\Lambda(n) = n\theta_1 + \varphi_1 - (2j+1)\pi$, where $j \in \mathbb{Z}$ is the unique integer such that $-\pi < \Lambda(n) \leq \pi$. We now have:

$$\begin{aligned} \frac{u_n}{\rho^n} &= 2|c_1| + 2|c_1| \cos(n\theta_1 + \varphi_1) + r(n) \\ &= 2|c_1|(1 + \cos(\Lambda(n) + \pi)) + r(n) \\ &\geq 2|c_1|g(\Lambda(n)) - (1 - \varepsilon)^n, \end{aligned}$$

where the inequality holds provided that $n > N$.

By Equation (4.12), when $|\Lambda(n)| \geq 1$, we have $\frac{u_n}{\rho^n} \geq \frac{11}{12}|c_1| - (1 - \varepsilon)^n$. It follows easily that $u_n/\rho^n \geq 0$ provided that $|\Lambda(n)| \geq 1$ and $n > N'$, for some $N' = 2^{\|\mathbf{u}\|^{O(1)}}$.

Recall that for $n > M$, $n\theta_1 + \varphi_1$ can never be an odd integer multiple of π , i.e., $\Lambda(n) \neq 0$. We now claim that there is an absolute constant $K \in \mathbb{N}$ such that, for $n > M$, we have $|\Lambda(n)| > n^{-\|\mathbf{u}\|^K}$.

To see this, write

$$\Lambda(n) = \frac{1}{i} \left(n \log \lambda_1 + \log \frac{c_1}{|c_1|} - (2j+1) \log(-1) \right).$$

In the above, if $c_1 \in \mathbb{R}$ and $c_1 > 0$, then simply remove the term $\log \frac{c_1}{|c_1|} = 0$ from the expression for $\Lambda(n)$, which would yield an even better lower bound than is obtained below. We may therefore assume without loss of generality that λ_1 and $c_1/|c_1|$ are different from 0 and 1.

Let $H \geq e$ be an upper bound for the heights of λ_1 and $c_1/|c_1|$, and let D be the largest of the degrees of λ_1 and $c_1/|c_1|$. Notice that the degree of $\mathbb{Q}(\lambda_1, c_1/|c_1|)$ over \mathbb{Q} is at most D^2 , and that $|j| \leq n$. We can thus invoke Theorem 3.4 to conclude that

$$\begin{aligned} |\Lambda(n)| &> \exp(-(48D^2)^{10} \log^2 H \log(2n+1)) \\ &= \frac{1}{(2n+1)^{(\log^2 H)(48D^2)^{10}}}, \end{aligned}$$

for $n > M$. The claim now follows by noting that both $\log H$ and D are bounded above by $\|\lambda_1\| + \|(c_1/|c_1|)\|$, and that the latter is in $\mathcal{O}(\|\mathbf{u}\|)$.

Thus when $|\Lambda(n)| < 1$ (and $n > M$), we have $g(\Lambda(n)) \geq g(n^{-\|\mathbf{u}\|^K})$. We can therefore find a polynomial $B \in \mathbb{Z}[x]$ such that

$$2|c_1|g(\Lambda(n)) \geq \frac{1}{B(n)},$$

requiring in addition that B have degree $\|\mathbf{u}\|^{O(1)}$ and height $2^{\|\mathbf{u}\|^{O(1)}}$, where the latter is achieved via bounds on the height of $|c_1|$. We can now invoke Proposition 3.1 to conclude that there is $N'' = 2^{\|\mathbf{u}\|^{O(1)}}$ such that $N'' \geq M$ and, for all $n > N''$, we have $\frac{1}{B(n)} > (1 - \varepsilon)^n$. Combining our various inequalities, we see that $u_n/\rho^n \geq 0$ provided that $n > \max\{N, N', N''\}$, which establishes ultimate positivity of \mathbf{u} and moreover once again provides an exponential bound on the index of possible violations of positivity, as required.

This concludes Subcase (i).

(ii) Finally, we turn to the situation in which the complex dominant roots γ_1 and $\overline{\gamma_1}$ are repeated. Using the same notation as above, we have

$$\begin{aligned} \frac{u_n}{\rho^n} &= a + (c_1 n + c) \lambda_1^n + (\overline{c_1} n + \overline{c}) \overline{\lambda_1}^n \\ &= a + n(c_1 \lambda_1^n + \overline{c_1} \overline{\lambda_1}^n) + c \lambda_1^n + \overline{c} \overline{\lambda_1}^n. \end{aligned}$$

Note that, unless $c_1 = 0$, the term $c_1 \lambda_1^n + \overline{c_1} \overline{\lambda_1}^n = 2|c_1| \cos(n\theta_1 + \varphi_1)$ is infinitely often negative and bounded away from zero, which immediately entails that \mathbf{u} can be neither positive nor ultimately positive. If $c_1 = 0$, on the other hand, we simply revert to an instance considered under Subcase (i).

4.3 Five Dominant Roots. If an LRS of order 5 has 5 distinct dominant roots, then each root is simple, and in Equation (4.5) we have that $m = 2$, $r(n)$ is identically 0, and the polynomials $A(n)$, $C_1(n)$, and $C_2(n)$ are all identically constant (cf. Section 2):

$$\frac{u_n}{\rho^n} = a + c_1 \lambda_1^n + \overline{c_1} \overline{\lambda_1}^n + c_2 \lambda_2^n + \overline{c_2} \overline{\lambda_2}^n,$$

for algebraic constants $a \in \mathbb{R}$ and $c_1, c_2 \in \mathbb{C}$.

Let $L = \{(v_1, v_2) \in \mathbb{Z}^2 : \lambda_1^{v_1} \lambda_2^{v_2} = 1\}$, and let B be a basis for L . Note that B can only have cardinality 0 (when L is trivial) or 1, since it is easily seen that the presence of two non-trivial independent integer multiplicative relationships over λ_1 and λ_2 would entail that λ_1 and λ_2 are roots of unity, contradicting the non-degeneracy of \mathbf{u} . Recall from Theorem 3.1 that the basis B can be computed in polynomial time, and moreover that elements of B may be assumed to have magnitude polynomial in $\|\mathbf{u}\|$.

If $B = \emptyset$, let

$$T = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| = |z_2| = 1\},$$

and if $B = \{(\ell_1, \ell_2)\}$, write

$$T = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| = |z_2| = 1 \text{ and } z_1^{\ell_1} z_2^{\ell_2} = 1\}.$$

Define $h : T \rightarrow \mathbb{R}$ by setting

$$h(z_1, z_2) = c_1 z_1 + \overline{c_1} \overline{z_1} + c_2 z_2 + \overline{c_2} \overline{z_2},$$

so that for all n , we have $u_n/\rho^n = a + h(\lambda_1^n, \lambda_2^n)$. By Corollary 3.1, the set $\{(\lambda_1^n, \lambda_2^n) : n \in \mathbb{N}\}$ is a dense subset of T . Since h is continuous, we immediately have that

$$\inf\{u_n/\rho^n : n \in \mathbb{N}\} = \min\{a + h(z_1, z_2) : (z_1, z_2) \in T\}.$$

It follows that \mathbf{u} is ultimately positive iff \mathbf{u} is positive iff $\min\{a + h(z_1, z_2) : (z_1, z_2) \in T\} \geq 0$ iff

$$(4.13) \quad \forall (z_1, z_2) \in T, a + h(z_1, z_2) \geq 0.$$

We now show how to rewrite Assertion (4.13) as a sentence in the first-order theory of the reals, i.e., involving only real-valued variables and first-order quantifiers, Boolean connectives, and integer constants together with the arithmetic operations of addition, subtraction, multiplication, and division.⁹ The idea is to separately represent the real and imaginary parts of each complex quantity appearing in Assertion (4.13), and combine them using real arithmetic so as to mimic the effect of complex arithmetic operations.

To this end, we use pairs of real variables x_1, y_1 and x_2, y_2 to represent z_1 and z_2 respectively: intuitively, $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$. Since the real constant a is algebraic, there is a formula $\sigma_a(x)$ which is true over the reals precisely for $x = a$. Likewise, the real and imaginary parts $\text{Re}(c_1)$, $\text{Im}(c_1)$, $\text{Re}(c_2)$, and $\text{Im}(c_2)$ of the complex algebraic constants c_1 and c_2 are themselves real algebraic, and can be represented as single-variable formulas in the first-order theory of the reals. All such formulas can readily be shown to have size polynomial in $\|\mathbf{u}\|$.

The terms $z_1^{\ell_1}$ and $z_2^{\ell_2}$ (if present) are simply expanded: for example, if ℓ_1 is positive, then $z_1^{\ell_1} = (x_1 + iy_1)^{\ell_1} = A_1(x_1) + iB_1(y_1)$, where A_1 and B_1 are polynomials with integer coefficients, and likewise for $z_2^{\ell_2}$. Note that since the magnitudes of ℓ_1 and ℓ_2 are polynomial in $\|\mathbf{u}\|$, so are $\|A_1\|$, $\|B_1\|$, $\|A_2\|$, and $\|B_2\|$. The case in which ℓ_1 or ℓ_2 is negative is handled similarly, with the additional use of a division operation.

Combining everything, we obtain a sentence τ of the first-order theory of the reals with division which is true iff Assertion (4.13) holds. τ makes use of at most 9 real variables: two for each of z_1 and z_2 , one for a , and one for each of $\text{Re}(c_1)$, $\text{Im}(c_1)$, $\text{Re}(c_2)$, $\text{Im}(c_2)$. In removing divisions from τ , the number of variables potentially increases to 11. Finally, the size of τ is polynomial in $\|\mathbf{u}\|$. We can therefore invoke Theorem 3.2 to conclude

⁹In Section 3, we did not include division as an allowable operation when we introduced the first-order theory of the reals; however instances of division can always be removed in linear time at the cost of introducing a linear number of existentially quantified fresh variables.

that Assertion (4.13) can be decided in time polynomial in $\|\mathbf{u}\|$.

This completes the proofs of Theorems 4.1 and 4.2.

5 Hardness at Order Six

Diophantine approximation is an old branch of number theory concerned, among other things, with problems related to approximating real numbers by rationals. It is a vast and active field of research with several hard, longstanding open problems. In this section, we present reductions from some of these open problems to questions of Positivity and Ultimate Positivity of integer LRS of order 6, and *a fortiori* of higher orders. In other words, we show that *if* Positivity or Ultimate Positivity are decidable for integer LRS of order 6, then certain hard open problems in Diophantine approximation become solvable.

We survey in cursory manner some of the key definitions and facts that are needed for our development. Results are stated largely without proofs—comprehensive reference works include [3, 50, 47].

For any real number x , the **Lagrange constant** (or **homogeneous Diophantine approximation constant**) $L_\infty(x)$ measures the extent to which x can be ‘well-approximated’ by rationals. It is defined as follows:

$$L_\infty(x) = \inf \left\{ c \in \mathbb{R} : \left| x - \frac{n}{m} \right| < \frac{c}{m^2} \text{ for infinitely many } n, m \in \mathbb{Z} \right\}.$$

Following Lagarias and Shallit’s terminology [33], we also define the (**homogeneous Diophantine approximation**) **type** of x :

$$L(x) = \inf \left\{ c \in \mathbb{R} : \left| x - \frac{n}{m} \right| < \frac{c}{m^2} \text{ for some } n, m \in \mathbb{Z} \right\}.$$

Khinchin showed in 1926 that almost all real numbers (in the measure-theoretic sense) have Lagrange constant and type equal to zero. Yet real numbers with non-zero Lagrange constant constitute an uncountable class known as the *badly approximable* numbers. The Lagrange constant and type of a real number x are closely linked to the continued fraction expansion of x , a fact which enabled Euler to prove that all algebraic numbers of degree 2 are badly approximable.

An old observation of Dirichlet shows that every real number has Lagrange constant at most 1. This bound was improved to $1/\sqrt{5}$ by Hurwitz in 1891, who also showed that it is achieved by the golden ratio. Markov proved in 1879 that every transcendental real number x has $L_\infty(x) \in [0, 1/3]$. Considerable further work has been devoted to the study of the Lagrange spectrum, which records the possible values taken on by Lagrange

constants—see, e.g., [24]. Despite this, nothing further is known about the Lagrange constant or type of the vast majority of transcendental numbers; for example, it is a longstanding open problem as to whether $L_\infty(\pi)$ is 0, $1/3$, or some value in between.

Let

$$\mathcal{A} = \{p + qi \in \mathbb{C} : p, q \in \mathbb{Q}, p^2 + q^2 = 1, \text{ and } p, q \neq 0\}$$

be the set of points on the unit circle in the complex plane with rational real and imaginary parts, excluding $\{1, -1, i, -i\}$. Note that this set is dense since $\frac{1-r^2}{1+r^2} + i\frac{2r}{1+r^2}$ always lies on the unit circle for any $r \in \mathbb{Q}$. Clearly, \mathcal{A} consists of algebraic numbers of degree 2, none of which is a root of unity: indeed, the primitive k th roots of unity are precisely the roots of the k th cyclotomic polynomial, whose degree is $\varphi(k)$, where φ is Euler’s totient function. Standard lower bounds on the latter imply that the only roots of unity of degree 2 are the 3rd, 4th, and 6th primitive roots of unity, all of which either have irrational imaginary part or are $\pm i$.

Write

$$\mathcal{T} = \left\{ \frac{\arg \alpha}{2\pi} : \alpha \in \mathcal{A} \right\}.$$

\mathcal{T} is a dense subset of $(-1/2, 1/2)$ consisting exclusively of transcendental numbers: indeed, for $t = \frac{\arg \alpha}{2\pi}$, we have $\alpha = e^{2\pi it} = (-1)^{2t}$. Since α is not a root of unity, t cannot be rational, and it follows that t must be transcendental by the Gelfond-Schneider theorem (see, e.g., [3]).

Recall that a real number x is **computable** if there is an algorithm which, given any rational $\varepsilon > 0$ as input, returns a rational q such that $|q - x| < \varepsilon$. We can now state our main hardness results:

THEOREM 5.1. *Suppose that Ultimate Positivity is decidable for integer LRS of order 6. Then, for any $t \in \mathcal{T}$, $L_\infty(t)$ is a computable number.*

THEOREM 5.2. *Suppose that Positivity is decidable for integer LRS of order 6. Then, for any $t \in \mathcal{T}$, $L(t)$ is a computable number.*

Theorems 5.1 and 5.2 strongly suggest that the decidability of Positivity and Ultimate Positivity for LRS of order 6 (and *a fortiori* higher orders) are unlikely to be achievable without major breakthroughs in analytic number theory. These theorems also have partial converses (which are omitted in the interest of brevity) which entail that, at least at order 6, proofs of *undecidability* would also have substantial implications regarding the Diophantine approximation of certain transcendental numbers.

We now proceed with the proof of both theorems.

Choose $p + qi \in \mathcal{A}$ and $r \in \mathbb{Q}$ such that $r > 0$. Let $\theta = \arg(p + qi)$, and write

$$\begin{aligned} u_n &= r \sin n\theta - n(1 - \cos n\theta) \\ v_n &= -r \sin n\theta - n(1 - \cos n\theta). \end{aligned}$$

It is not hard to see that $\mathbf{u} = \langle u_n \rangle_{n=0}^\infty$ and $\mathbf{v} = \langle v_n \rangle_{n=0}^\infty$ are both rational LRS of order 6. Indeed, writing $\lambda = p + qi$, both \mathbf{u} and \mathbf{v} are LRS with characteristic roots 1, λ , and $\bar{\lambda}$, each of which has multiplicity 2. The exponential polynomial solution for \mathbf{u} is

$$u_n = -n1^n + \frac{1}{2}(n - ri)\lambda^n + \frac{1}{2}(n + ri)\bar{\lambda}^n,$$

from which the order-6 recurrence relation can easily be extracted. Note that since λ and $\bar{\lambda}$ have rational real and imaginary parts, by induction u_n is rational for all $n \geq 0$. Naturally, a similar exercise can be carried out for \mathbf{v} .

For $n \geq 0$, let

$$w_n = \max\{u_n, v_n\} = r|\sin n\theta| - n(1 - \cos n\theta).$$

Given $\varepsilon \in (0, 1)$, there exists $\delta > 0$ such that, for all $x \in [-\delta, \delta]$, we have

$$(5.14) \quad (1 - \varepsilon)|x| \leq |\sin x| \leq |x|, \text{ and}$$

$$(5.15) \quad (1 - \varepsilon)\frac{x^2}{2} \leq 1 - \cos x \leq \frac{x^2}{2}$$

Moreover, there exists $N \in \mathbb{N}$ with $2r/N \leq \delta$ such that, for all $x \in (-\pi, \pi]$,

$$(5.16) \quad \text{if } 1 - \cos x < \frac{2r}{N}, \text{ then } |x| \leq \delta.$$

For $x \in \mathbb{R}$, recall that $[x]_{2\pi}$ denotes the distance from x to the closest integer multiple of 2π . Let $t = \theta/2\pi$. It is straightforward to show that

$$(5.17) \quad 2\pi L_\infty(t) = \liminf_{m \in \mathbb{N}} m[m(2\pi t)]_{2\pi} = \liminf_{m \in \mathbb{N}} m[m\theta]_{2\pi}$$

and

$$(5.18) \quad 2\pi L(t) = \inf_{m \in \mathbb{N}} m[m(2\pi t)]_{2\pi} = \inf_{m \in \mathbb{N}} m[m\theta]_{2\pi}.$$

We now assert the following:

Claim 1: For any $m \geq N$, if $w_m > 0$, then $m[m\theta]_{2\pi} < \frac{2r}{1 - \varepsilon}$.

Claim 2: For any $m \geq N$, if $m[m\theta]_{2\pi} < 2r(1 - \varepsilon)$, then $w_m > 0$.

To prove Claim 1, assume that $m \geq N$ and $w_m > 0$. Then:

$$\begin{aligned} m(1 - \cos m\theta) &< r && \text{[by definition of } w_m\text{]} \\ \Rightarrow 1 - \cos m\theta &< \frac{r}{m} < \frac{2r}{m} \leq \frac{2r}{N} \\ \Rightarrow [m\theta]_{2\pi} &\leq \delta && \text{[by (5.16)]} \\ \Rightarrow 0 < w_m &&& \\ &\leq r[m\theta]_{2\pi} - m(1 - \varepsilon)\frac{([m\theta]_{2\pi})^2}{2} && \text{[(5.14), (5.15)]} \\ \Rightarrow m[m\theta]_{2\pi} &< \frac{2r}{1 - \varepsilon}, \end{aligned}$$

as required.

For Claim 2, assume that $m \geq N$ and $m[m\theta]_{2\pi} < 2r(1 - \varepsilon)$. Then $[m\theta]_{2\pi} \leq 2r/N \leq \delta$, whence $w_n \geq r(1 - \varepsilon)[m\theta]_{2\pi} - m\frac{([m\theta]_{2\pi})^2}{2}$ by (5.14) and (5.15), and also $m\frac{([m\theta]_{2\pi})^2}{2} < r(1 - \varepsilon)[m\theta]_{2\pi}$.¹⁰ Combining the last two inequalities yields $w_n > 0$ as required.

Observe that if $-\mathbf{u}$ and $-\mathbf{v}$ are both ultimately positive,¹¹ then for all sufficiently large m , we have $w_m \leq 0$, and therefore, by Claim 2, $m[m\theta]_{2\pi} \geq 2r(1 - \varepsilon)$. Since this holds for all $\varepsilon \in (0, 1)$, it follows from Equation (5.17) that $L_\infty(t) \geq r/\pi$.

On the other hand, if one or both of $-\mathbf{u}$ and $-\mathbf{v}$ fail to be ultimately positive, then there must be infinitely many values of m such that $w_m > 0$. Claim 1 and Equation (5.17) then entail that $L_\infty(t) \leq r/\pi$.

Since r can be chosen arbitrarily, this establishes Theorem 5.1.

A similar procedure can be used to approximate $L(t)$. Note that arbitrarily good upper bounds can always be guessed and, if correct, be verified effectively, by enumerating pairs of integers until a suitable pair is found.¹²

Suppose now that we wish to validate a purported lower bound $b < L(t)$. Guess rational values of r and ε such that $2\pi b < 2r(1 - \varepsilon) < \frac{2r}{1 - \varepsilon} < 2\pi L(t)$. Note that one can readily compute the value of the corresponding integer N in the notation of our proof. Invoke the Positivity oracle on the LRS $\langle -u_m \rangle_{m=N}^\infty$ and $\langle -v_m \rangle_{m=N}^\infty$. The outcome must be that both are positive, otherwise there would be some value of $m \geq N$ such that $w_m > 0$, from which we would conclude

¹⁰Recall that $p + qi$ is not a root of unity, and hence $[m\theta]_{2\pi} \neq 0$.

¹¹Recall from Section 4 that decision procedures for Positivity and Ultimate Positivity of integer LRS are readily applicable to rational LRS.

¹²Note that this requires some numerical analysis, which we take for granted, in order to perform approximations with sufficient precision.

via Claim 1 that $m[m\theta]_{2\pi} < \frac{2r}{1-\varepsilon}$, contradicting our assumption that $\frac{2r}{1-\varepsilon} < 2\pi L(t)$.

Since both LRS are revealed to be positive, we know that for all $m \geq N$, $w_m \leq 0$ and therefore (thanks to Claim 2) that $m[m\theta]_{2\pi} \geq 2r(1-\varepsilon)$. It now suffices to verify individually each value of $m \in \{0, \dots, N-1\}$ to conclude that $2\pi L(t) \geq 2r(1-\varepsilon) > 2\pi b$, as required. This completes the proof of Theorem 5.2.

Let us finally remark that other hardness results, similar in both form and spirit to Theorems 5.1 and 5.2, can also be formulated, notably via the use of techniques on *inhomogeneous* Diophantine approximation of certain transcendental numbers.

6 Extensions and Future Work

Several of the results presented in this paper have natural extensions or generalisations, some of which we briefly mention here.

Define an LRS $\mathbf{u} = \langle u_n \rangle_{n=0}^\infty$ to be **strictly positive** (respectively **ultimately strictly positive**) if $u_n > 0$ for all n (respectively for all sufficiently large n). An examination of our proofs readily shows that all our decidability and complexity results, with the exception of the decidability and complexity of Positivity for integer LRS of order 5, carry over without difficulty to the analogous strict formulation. A useful observation in this regard is that for non-degenerate LRS, Ultimate Positivity and Ultimate Strict Positivity agree: indeed, as can be seen from the proof of the Skolem-Mahler-Lech theorem [26], any non-degenerate LRS is either identically zero or has only finitely many zeros. Let us also mention that our Diophantine-approximation hardness results are easily seen to carry over *mutatis mutandis* to Strict Positivity and Ultimate Strict Positivity.

All our decidability results also carry over to LRS over real algebraic numbers, as can readily be seen by examining the relevant proofs. Our complexity upper bounds, however, are more delicate, and it is an open question whether they continue to hold in the algebraic setting. Hardness results, on the other hand, obviously carry over to the more general algebraic world.

It seems likely that the techniques developed in this paper could be usefully deployed to tackle other natural decision problems for linear recurrence sequences, such as divergence to infinity, reachability and ultimate reachability of semi-linear sets, etc. In turn such decision procedures—or corresponding hardness results—may find applications in some of the areas mentioned in the Introduction, such as the analysis of termination of linear programs or the behaviour of discrete linear dynamical systems. More ambitiously, in the spirit of synthesis, one could seek to explore computational problems for *parametric* LRS, where the aim is to char-

acterise ranges for the parameters guaranteeing certain behaviours, etc.

Another interesting question concerns the complexity of Positivity at low orders. Recall that the PosSLP oracle used in our main decision procedure is invoked to check whether the quantity $\vec{v}^T M^n \vec{w}$ is strictly negative, where M is a $k \times k$ matrix of integers, \vec{v} and \vec{w} are k -dimensional integer vectors, and n is encoded in binary. It is conceivable—especially for small fixed k , as in the situation at hand—that the complexity of this problem is significantly lower than that of PosSLP. See [31] for initial progress on related questions.

Finally, the various discrete problems discussed in the present paper also have natural counterparts in a continuous setting. See [9], for example, which studies the Skolem and Positivity Problems over continuous time using similar tools. This remains a largely unexplored research landscape.

References

- [1] M. Agrawal, S. Akshay, B. Genest, and P. S. Thiagarajan. Approximate verification of the symbolic dynamics of Markov chains. In *Proc. Symp. on Logic in Comp. Sci. (LICS)*. IEEE, 2012.
- [2] E. Allender, P. Bürgisser, J. Kjeldgaard-Pedersen, and P. B. Miltersen. On the complexity of numerical analysis. *SIAM J. Comput.*, 38(5), 2009.
- [3] A. Baker. *Transcendental Number Theory*. Cambridge University Press, 1975.
- [4] A. Baker and G. Wüstholz. Logarithmic forms and group varieties. *Jour. Reine Angew. Math.*, 442, 1993.
- [5] S. Basu, R. Pollack, and M.-F. Roy. *Algorithms in Real Algebraic Geometry*. Springer, 2nd edition, 2006.
- [6] W. J. Baumol. *Economic Dynamics*. Prentice Hall, 3rd edition, 1970.
- [7] D. Beauquier, A. M. Rabinovich, and A. Slissenko. A logic of probability with decidable model checking. *J. Log. Comput.*, 16(4), 2006.
- [8] J. P. Bell and S. Gerhold. On the positivity set of a linear recurrence. *Israel Jour. Math.*, 57, 2007.
- [9] P. C. Bell, J.-C. Delvenne, R. M. Jungers, and V. D. Blondel. The continuous Skolem-Pisot problem. *Theor. Comput. Sci.*, 411(40-42), 2010.
- [10] A. M. Ben-Amram. Mortality of iterated piecewise affine functions over the integers: Decidability and complexity. In *Proc. Intern. Symp. on Theoret. Aspects of Comp. Sci. (STACS)*, volume 20 of *LIPICs*. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2013.
- [11] A. M. Ben-Amram, S. Genaim, and A. N. Masud. On the termination of integer loops. *ACM Trans. Program. Lang. Syst.*, 34(4), 2012.
- [12] J. Berstel and M. Mignotte. Deux propriétés décidables des suites récurrentes linéaires. *Bull. Soc. Math. France*, 104, 1976.

- [13] V. D. Blondel, E. Jeandel, P. Koiran, and N. Portier. Decidable and undecidable problems about quantum automata. *SIAM J. Comput.*, 34(6), 2005.
- [14] V. D. Blondel and N. Portier. The presence of a zero in an integer linear recurrent sequence is NP-hard to decide. *Linear Algebra and Its Applications*, 351–352, 2002.
- [15] N. Bourbaki. *Elements of Mathematics: General Topology (Part 2)*. Addison-Wesley, 1966.
- [16] M. Bozga, R. Iosif, and Filip Konečný. Deciding conditional termination. In *Proc. Intern. Conf. on Tools and Algorithms for the Construction and Analysis of Systems (TACAS)*, volume 7214 of *LNCS*. Springer, 2012.
- [17] M. Braverman. Termination of integer linear programs. In *Proc. Intern. Conf. on Computer Aided Verification (CAV)*, volume 4144 of *LNCS*. Springer, 2006.
- [18] J. R. Burke and W. A. Webb. Asymptotic behavior of linear recurrences. *Fib. Quart.*, 19(4), 1981.
- [19] J.-Y. Cai, R. J. Lipton, and Y. Zalcstein. The complexity of the A B C problem. *SIAM J. Comput.*, 29(6), 2000.
- [20] V. Chonev, J. Ouaknine, and J. Worrell. The Orbit Problem in higher dimensions. In *Proc. Symp. on the Theory of Computing (STOC)*. ACM, 2013.
- [21] H. Cohen. *A Course in Computational Algebraic Number Theory*. Springer-Verlag, 1993.
- [22] G. E. Collins. Quantifier elimination for real closed fields by cylindrical algebraic decomposition. In *Proc. 2nd GI Conf. Automata Theory and Formal Languages*. Springer-Verlag, 1975.
- [23] B. Cook, A. Podelski, and A. Rybalchenko. Proving program termination. *Commun. ACM*, 54(5), 2011.
- [24] T. W. Cusick and M. E. Flahive. *The Markoff and Lagrange Spectra*. American Mathematical Society, 1989.
- [25] H. Derksen, E. Jeandel, and P. Koiran. Quantum automata and algebraic groups. *J. Symb. Comput.*, 39(3-4), 2005.
- [26] G. Everest, A. van der Poorten, I. Shparlinski, and T. Ward. *Recurrence Sequences*. American Mathematical Society, 2003.
- [27] G. Ge. *Algorithms Related to Multiplicative Representations of Algebraic Numbers*. PhD thesis, U.C. Berkeley, 1993.
- [28] I. Gyori and G. Ladas. *Oscillation Theory of Delay Differential Equations*. Oxford Mathematical Monographs. Oxford University Press, 1991.
- [29] V. Halava, T. Harju, and M. Hirvensalo. Positivity of second order linear recurrent sequences. *Discrete Applied Mathematics*, 154(3), 2006.
- [30] V. Halava, T. Harju, M. Hirvensalo, and J. Karhumäki. Skolem’s problem — on the border between decidability and undecidability. Technical Report 683, Turku Centre for Computer Science, 2005.
- [31] M. Hirvensalo, J. Karhumäki, and A. Rabinovich. Computing partial information out of intractable: Powers of algebraic numbers as an example. *Jour. Number Theory*, 130, 2010.
- [32] R. Kannan and R. J. Lipton. Polynomial-time algorithm for the orbit problem. *Jour. ACM*, 33(4), 1986.
- [33] J. C. Lagarias and J. O. Shallit. Linear fractional transformations of continued fractions with bounded partial quotients. *Journal de Théorie des Nombres de Bordeaux*, 9, 1997.
- [34] V. Laohakosol. Personal communication, July 2013.
- [35] V. Laohakosol and P. Tangsupphathawat. Positivity of third order linear recurrence sequences. *Discrete Applied Mathematics*, 157(15), 2009.
- [36] A. Lindenmayer and G. Rozenberg, editors. *Automata, Languages, Development*. North-Holland, 1976.
- [37] R. J. Lipton. Mathematical embarrassments. *Blog entry*, December 2009. <http://rjlipton.wordpress.com/2009/12/26/mathematical-embarrassments/>.
- [38] L. L. Liu. Positivity of three-term recurrence sequences. *Electr. J. Comb.*, 17(1), 2010.
- [39] D. W. Masser. Linear relations on algebraic groups. In *New Advances in Transcendence Theory*. Cambridge University Press, 1988.
- [40] M. Mignotte. Some useful bounds. In *Computer Algebra*, 1982.
- [41] M. Mignotte, T. N. Shorey, and R. Tijdeman. The distance between terms of an algebraic recurrence sequence. *Journal für die reine und angewandte Mathematik*, 349, 1984.
- [42] K. Nagasaka and J.-S. Shiue. Asymptotic positiveness of linear recurrence sequences. *Fib. Quart.*, 28(4), 1990.
- [43] J. Ouaknine and J. Worrell. Decision problems for linear recurrence sequences. In *Proc. Intern. Workshop on Reachability Problems (RP)*, volume 7550 of *LNCS*. Springer, 2012.
- [44] V. Pan. Optimal and nearly optimal algorithms for approximating polynomial zeros. *Computers & Mathematics with Applications*, 31(12), 1996.
- [45] A. Podelski and A. Rybalchenko. A complete method for the synthesis of linear ranking functions. In *Proc. Intern. Conf. on Verif., Model Checking, and Abstract Interpretation (VMCAI)*, volume 2937 of *LNCS*. Springer, 2004.
- [46] J. Renegar. On the computational complexity and geometry of the first-order theory of the reals. Part I: Introduction. Preliminaries. The geometry of semi-algebraic sets. The decision problem for the existential theory of the reals. *J. Symb. Comp.*, 1992.
- [47] A. M. Rockett and P. Szűsz. *Continued Fractions*. World Scientific, 1992.
- [48] G. Rozenberg and A. Salomaa. *Cornerstones of Undecidability*. Prentice Hall, 1994.
- [49] A. Salomaa. Growth functions of Lindenmayer systems: Some new approaches. In A. Lindenmayer and G. Rozenberg, editors, *Automata, Languages, Development*. North-Holland, 1976.
- [50] W. M. Schmidt. Diophantine approximation. In *Lect. Notes in Math.*, volume 785, 1980.

- [51] M. Soittola. On D0L synthesis problem. In A. Lindenmayer and G. Rozenberg, editors, *Automata, Languages, Development*. North-Holland, 1976.
- [52] P. Tangsupphathawat, N. Punnim, and V. Laohakosol. The positivity problem for fourth order linear recurrence sequences is decidable. *Colloq. Math.*, 128(1), 2012.
- [53] T. Tao. Open question: effective Skolem-Mahler-Lech theorem. *Blog entry*, May 2007. <http://terrytao.wordpress.com/2007/05/25/open-question-effective-skolem-mahler-lech-theorem/>.
- [54] S. P. Tarasov and M. N. Vyalyi. Orbits of linear maps and regular languages. In *Proc. Intern. Comp. Sci. Symp. in Russia (CSR)*, volume 6651 of *LNCS*. Springer, 2011.
- [55] A. Tarski. *A Decision Method for Elementary Algebra and Geometry*. University of California Press, 1951.
- [56] A. Tiwari. Termination of linear programs. In *Proc. Intern. Conf. on Comp. Aided Verif. (CAV)*, volume 3114 of *LNCS*. Springer, 2004.
- [57] N. K. Vereshchagin. The problem of appearance of a zero in a linear recurrence sequence (in Russian). *Mat. Zametki*, 38(2), 1985.
- [58] K. Yokoyama, Z. Li, and I. Nemes. Finding roots of unity among quotients of the roots of an integral polynomial. In *Proc. Intern. Symp. on Symb. and Algebraic Comp.*, 1995.