

Effective Divergence Analysis for Linear Recurrence Sequences

Shaul Almagor

Department of Computer Science, Oxford University, UK
shaull.almagor@cs.ox.ac.uk

Brynmor Chapman

MIT CSAIL
brynmor@mit.edu

Mehran Hosseini

Department of Computer Science, Oxford University, UK
mehran.hosseini@cs.ox.ac.uk

Joël Ouaknine¹

Max Planck Institute for Software Systems, Germany &
Department of Computer Science, Oxford University, UK
joel@mpi-sws.org

James Worrell²

Department of Computer Science, Oxford University, UK
jbw@cs.ox.ac.uk

Abstract

We study the growth behaviour of rational linear recurrence sequences. We show that for low-order sequences, divergence is decidable in polynomial time. We also exhibit a polynomial-time algorithm which takes as input a divergent rational linear recurrence sequence and computes effective fine-grained lower bounds on the growth rate of the sequence.

2012 ACM Subject Classification Computing methodologies → Algebraic algorithms, Theory of computation → Logic and verification

Keywords and phrases Linear recurrence sequences, Divergence, Algebraic numbers, Positivity

Digital Object Identifier 10.4230/LIPIcs.CONCUR.2018.42

1 Introduction

Linear recurrence sequences (LRS), such as the Fibonacci numbers, permeate a wide range of scientific fields, from economics and theoretical biology to computer science and mathematics. In computer-aided verification, for example, LRS techniques play a key rôle in the termination analysis of a large class of simple while loops—see [19] for a short survey on this topic. Likewise, the ergodic behaviour of Markov chains in probability theory [1], or the stability of supply-and-demand price equilibria in laggy markets in economics (the so-called ‘cobweb model’) [4] can be analysed through an examination of the asymptotic behaviour of certain types of LRS; in particular, instability of price equilibria corresponds precisely to *divergence* of the associated LRS.

¹ Supported by ERC grant AVS-ISS (648701)

² Supported by EPSRC Fellowship EP/N008197/1



In this paper, we undertake a systematic and fine-grained analysis of the growth behaviour of rational linear recurrence sequences from the point of view of effectiveness and complexity. In order to describe our main results, we first require some preliminary definitions. A sequence of real numbers $\mathbf{u} = \langle u_n \rangle_{n=1}^\infty$ is said to satisfy a *linear recurrence of order k* if there are real numbers a_1, \dots, a_{k+1} such that

$$u_{n+k} = a_1 u_{n+k-1} + \dots + a_{k-1} u_{n+1} + a_k u_n + a_{k+1} \quad (1)$$

for all $n \in \mathbb{N}$. Such a recurrence is said to be *homogeneous* if $a_{k+1} = 0$ and *inhomogeneous* if $a_{k+1} \neq 0$. The *characteristic polynomial* of the recurrence is

$$p(x) := x^k - a_1 x^{k-1} - \dots - a_{k-1} x - a_k.$$

The zeros of p are called the *characteristic roots*. A characteristic root of maximum modulus is said to be *dominant* and its modulus is the *dominant modulus*. The *multiplicity* of a characteristic root γ is the maximal $m \in \mathbb{N}$ such that $(x - \gamma)^m$ divides $p(x)$.

An LRS is said to be *rational* if it consists of rational numbers, *integral* if it consists of integers, and *algebraic* if it consists of algebraic numbers. An LRS is *simple* if all of its characteristic roots have multiplicity 1, and is *non-degenerate* if no ratio of two distinct characteristic roots is a root of unity.³

We say that an LRS \mathbf{u} *diverges to ∞* if $\lim_{n \rightarrow \infty} u_n = \infty$ (technically speaking: for all $T \in \mathbb{N}$, there exists $N \in \mathbb{N}$ such that, for all $n \geq N$, we have $u_n \geq T$). We also say that \mathbf{u} is *absolutely divergent* (or *diverges in absolute value*) if $\lim_{n \rightarrow \infty} |u_n| = \infty$.

The LRS \mathbf{u} is said to be *positive* if $u_n \geq 0$ for all $n \geq 1$, and *ultimately positive* if there is some $N \in \mathbb{N}$ such that $u_n \geq 0$ for all $n \geq N$.

A celebrated result from the 1930s, the Skolem-Mahler-Lech theorem (see [10]), implies that all non-degenerate integral LRS are absolutely divergent. This statement is however *non-effective* in a very basic sense: given a finite representation of a non-degenerate integral LRS \mathbf{u} , there is no known algorithm to compute a bound N such that $u_n \neq 0$ for $n \geq N$. It is also worth pointing out that the divergence assertion fails in general for non-integral LRS.

The question of the so-called *rate of absolute divergence* for non-degenerate integral LRS was subsequently extensively studied; see [10, Sec. 2.4] for an account of some of the key results accumulated over the last several decades. To begin with, a fairly straightforward fact is the following: if \mathbf{u} is an algebraic LRS of order k with dominant modulus ρ , then there is an effectively computable constant a such that, for all $n \geq 1$, $|u_n| \leq a\rho^n n^k$. In the 1970s, a conjecture was formulated to the effect that any non-degenerate integral LRS has, essentially, the maximal possible growth rate (see the next theorem for a precise statement). The conjecture was finally settled positively independently by Evertse [11] and by van der Poorten and Schlickewei [23]:

► **Theorem 1.** *For any non-degenerate algebraic LRS \mathbf{u} of dominant modulus $\rho > 1$, and any $\varepsilon > 0$, there exists a constant N such that, for all $n \geq N$, we have $|u_n| \geq \rho^{(1-\varepsilon)n}$.*

This is a highly non-trivial result making use of deep number-theoretic tools concerning bounds on the sum of S -units. Unfortunately, the proof is not effective, in the sense that given $\varepsilon > 0$, it does not provide estimates for the corresponding value of N . This effectiveness

³ For most practical purposes—and certainly for all of the computational tasks considered in this paper—LRS can be assumed to be non-degenerate, since any degenerate LRS can be effectively decomposed into a finite number of non-degenerate LRS; moreover this reduction can be carried out in polynomial time for rational LRS of bounded order [10, 16].

issue is described as “an important open problem” in [10], where it is furthermore suggested that any progress on the matter would likely hinge upon substantial improvements of deep number-theoretic results, such as Roth’s theorem, the prospects of which currently appear to be remote.

Nevertheless—and in particular for algorithmic applications in computer science—effectiveness is of central importance. The sharpest known results in this vein are due to Mignotte [13] as well as Shorey and Stewart [21], capping a long line of work in this area:

► **Theorem 2.** *For any homogeneous non-degenerate integral LRS \mathbf{u} of order at most 3 with dominant modulus ρ , there are effective constants a and d such that, for all $n \geq 1$, we have $|u_n| \geq \frac{a\rho^n}{n^d}$.*

For most problems in computer science and automated verification, such as the analysis of the long-run behaviour of dynamical systems or the termination of linear while loops, the primary notion of *divergence* is clearly much more relevant than that of ‘divergence in absolute value’. In view of the above results, however, one might expect that little could be said about effective rates of divergence. Somewhat surprisingly, divergence does turn out to be significantly more tractable than absolute divergence. At a high level, the main results of this paper can now be summarised as follows:

Given a rational LRS \mathbf{u} , homogeneous or inhomogeneous, either of order at most 5, or, if the LRS is simple, of order at most 8, we can carry out the following tasks in polynomial time:

- decide if \mathbf{u} diverges to ∞ or not; and
- in divergent instances, provide effective fine-grained lower bounds on the rate of divergence of \mathbf{u} .

The precise statements can be found in Theorems 12 and 13. The most obvious contrast in comparison with Theorem 2 is the higher order of LRS that can be handled effectively (5 and 8 versus 3). It is also worth noting, however, that our results apply more generally to rational (as opposed to integral) LRS, and that we can handle inhomogeneous sequences at no cost—this is remarkable in that the folklore wisdom usually broadly equates inhomogeneous LRS of order k with homogeneous LRS of order $k + 1$ (this assertion, as well as the manner in which we circumvent it, are made precise in the main body of the paper).

Finally, let us point out that our analysis of divergence rates relies, among others, on improvements to results concerning the positivity and ultimate positivity of LRS, which were originally developed in [17, 16, 18]. As a by-product, therefore, stronger results on the Positivity and Ultimate Positivity Problems—notably dealing with inhomogeneous LRS—can be found in the present paper, in particular in the form of Theorems 19 and 20.

2 Preliminaries

2.1 Linear Recurrence Sequences

Let us start by reformulating the notion of linear recurrence more abstractly as follows. Define the *shift operator* $E : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ by $E(f)(n) = f(n + 1)$ for a sequence $f \in \mathbb{R}^{\mathbb{N}}$. The polynomial ring $\mathbb{R}[E]$ acts on the set of sequences $\mathbb{R}^{\mathbb{N}}$ on the left in a natural way, turning $\mathbb{R}^{\mathbb{N}}$ into a left $\mathbb{R}[E]$ module. Then a sequence $\mathbf{u} = \langle u \rangle_{n=1}^{\infty}$ satisfies the recurrence equation (1) if and only if $p(E) \cdot \mathbf{u} = a_{k+1} \cdot \mathbf{1}$, where p is the characteristic polynomial of the recurrence and $\mathbf{1}$ is the all-ones sequence.

The following homogenization construction is well known.

► **Proposition 3.** *Let $\mathbf{u} = \langle u_n \rangle_{n=1}^\infty$ satisfy an inhomogeneous linear recurrence of order k . Then \mathbf{u} satisfies a homogeneous recurrence of order $k + 1$.*

Proof. By assumption we have that $p(E) \cdot \mathbf{u} = \mathbf{c}$ for some monic polynomial $p(x)$ of degree k and constant sequence \mathbf{c} . Writing $q(x) = (x - 1)p(x)$, we have $q(E) \cdot \mathbf{u} = (E - 1) \cdot \mathbf{c} = 0$. ◀

We have the following partial converse to Proposition 3.

► **Proposition 4.** *Let $\mathbf{u} = \langle u_n \rangle_{n=1}^\infty$ satisfy a homogeneous linear recurrence of order $k + 1$ with a positive real characteristic root ρ . Then the sequence $\mathbf{v} = \langle v_n \rangle_{n=1}^\infty$ defined by $v_n = \frac{u_n}{\rho^n}$ satisfies an inhomogeneous linear recurrence of order k .*

Proof. By assumption, \mathbf{u} satisfies the recurrence $f(E) \cdot \mathbf{u} = 0$ for some monic polynomial $f(x) \in \mathbb{R}[x]$ of degree $k + 1$ that has a positive real root ρ . Define a sequence $\mathbf{v} = \langle v_n \rangle_{n=1}^\infty$ by $v_n := \frac{u_n}{\rho^n}$ for all $n \in \mathbb{N}$. Then \mathbf{v} satisfies the recurrence $g(E) \cdot \mathbf{v} = 0$ where g is the monic polynomial $g(x) = \rho^{-(k+1)} f(\rho x)$.

But $g(1) = 0$ and hence we have the factorization $g(x) = (x - 1)h(x)$ for some monic polynomial $h(x) \in \mathbb{R}[x]$. It follows that $(E - 1)h(E) \cdot \mathbf{v} = 0$ and hence $h(E) \cdot \mathbf{v}$ is constant, i.e., \mathbf{v} satisfies an inhomogeneous recurrence of order k . ◀

Let $\|\mathbf{u}\|$ denote the binary representation length⁴ of \mathbf{u} . We remark that the transformations back and forth between homogeneous and inhomogeneous LRS can be carried out in polynomial time in $\|\mathbf{u}\|$ if the given LRS have real algebraic coefficients. For an inhomogeneous LRS \mathbf{u} of order k , we refer to the corresponding homogeneous LRS obtained as per Proposition 3 as the *homogenization* of \mathbf{u} , denoted $\text{HOM}(\mathbf{u})$. The proof of Proposition 3 gives us the following useful property.

► **Property 5.** *The characteristic roots of $\text{HOM}(\mathbf{v})$ are the same as those of \mathbf{v} , with the same multiplicities, except for the characteristic root 1, which always occurs in $\text{HOM}(\mathbf{v})$, and whose multiplicity is $m + 1$, where m is the multiplicity of 1 in \mathbf{v} .*

Consider an LRS \mathbf{u} with integer coefficients. Then, since the characteristic polynomial p of an LRS \mathbf{u} has integer coefficients, the characteristic roots of \mathbf{u} comprise real-algebraic roots $\{\rho_1, \dots, \rho_d\}$, and conjugate pairs of complex-algebraic roots $\{\gamma_1, \overline{\gamma_1}, \dots, \gamma_m, \overline{\gamma_m}\}$. There are now univariate polynomials $A_1, \dots, A_d \in \mathbb{R}[x]$ with real-algebraic coefficients and C_1, \dots, C_m with complex-algebraic coefficients such that, for every $n \geq 0$,

$$u_n = \sum_{i=1}^d A_i(n)\rho_i^n + \sum_{j=1}^m (C_j(n)\gamma_j^n + \overline{C_j(n)}\overline{\gamma_j}^n)$$

This expression is referred to as the *exponential polynomial* solution of \mathbf{u} . The degree of each of the polynomials is strictly smaller than the multiplicity of the corresponding root. For a fixed order k , the coefficients appearing in the polynomials can be computed in time polynomial in $\|\mathbf{u}\|$.

We now turn to present two results regarding the asymptotic analysis of LRS.

The following result due to Braverman [6] enables us to reason about the complex part of the exponential polynomial above.

⁴ in general, we denote by $\|\cdot\|$ the binary-representation length of objects.

► **Lemma 6** (Complex Units Lemma). *Let $\zeta_1, \zeta_2, \dots, \zeta_m \in \mathbf{S}_1 \setminus \{1\}$ be distinct complex numbers (where $\mathbf{S}_1 = \{z \in \mathbb{C} : |z| = 1\}$), and let $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{C} \setminus \{0\}$. Set $z_n := \sum_{k=1}^m \alpha_k \zeta_k^n$. Then there exists $c < 0$ such that for infinitely many n , $\operatorname{Re}(z_n) < c$.*

In particular, Lemma 6 immediately implies that an LRS without a real dominant characteristic root, is neither positive, ultimately positive, nor divergent.

Finally, the following proposition from [16] allows us to bound the growth rate of the low-order terms in the exponential polynomial of an LRS.

► **Proposition 7.** *Consider an LRS $\mathbf{u} = \langle u_n \rangle_{n=1}^\infty$ of bounded order, with dominant modulus ρ , and write*

$$\frac{u_n}{\rho^n} = A(n) + \sum_{i=1}^m \left(C_i(n) \lambda_i^n + \overline{C_i(n)} \overline{\lambda_i}^n \right) + r(n)$$

where A is a real polynomial, C_i are non-zero complex polynomials, $\rho \lambda_i$ and $\rho \overline{\lambda_i}$ are conjugate pairs of non-real dominant roots of \mathbf{u} , and r is an exponentially decaying function.

We can compute in polynomial time $\epsilon \in (0, 1)$ and $N \in \mathbb{N}$ such that

$$\frac{1}{\epsilon} = 2^{\|\mathbf{u}\|^{O(1)}},$$

$$N = 2^{\|\mathbf{u}\|^{O(1)}},$$

for all $n > N$, $|r(n)| < (1 - \epsilon)^n$.

2.2 Mathematical Tools

In this section we introduce several tools that will be used throughout the paper.

Algebraic Numbers A complex number α is *algebraic* if it is a root of a polynomial $p \in \mathbb{Z}[x]$. The *defining polynomial* of α is the unique monic polynomial of minimal degree that has α as a root, and is denoted p_α . The *degree* and the *height* of α are the degree and the height (i.e., maximum absolute value of the coefficients) of p_α , respectively. An algebraic number α can be represented by a polynomial that has α as a root, along with an approximation of α by a complex number with rational real and imaginary parts. We denote by $\|\alpha\|$ the representation length of α . Basic arithmetic operations as well as equality testing and comparisons for algebraic numbers can be carried out in polynomial time (see [5, 8] for efficient algorithms).

The following lemma from [17] is a consequence of the celebrated lower bound for linear forms in logarithms due to Baker and Wüstholz [3].

► **Lemma 8.** *There exists $D \in \mathbb{N}$ such that, for all algebraic numbers $\lambda, \zeta \in \mathbb{C}$ of modulus 1, and for all $n \geq 2$, if $\lambda^n \neq \zeta$, then $|\lambda^n - \zeta| > \frac{1}{n^{(\|\lambda\| + \|\zeta\|)^D}}$.*

Multiplicative Relations Multiplicative relations between characteristic roots of an LRS play a key role in our analysis. The following result, due to Masser [12] enables us to efficiently elicit these relationships.

► **Theorem 9.** *Let m be fixed, and let $\lambda_1, \dots, \lambda_m$ be complex algebraic numbers of modulus 1. Let $L = \{(v_1, \dots, v_m) \in \mathbb{Z}^m : \lambda_1^{v_1} \dots \lambda_m^{v_m} = 1\}$ be the group of multiplicative relations between the λ_i . L has a basis $\{\ell_1, \dots, \ell_p\} \subseteq \mathbb{Z}^m$ (with $p \leq m$), where the entries of each of the ℓ_j are all polynomially bounded in $\|\lambda_1\| + \dots + \|\lambda_m\|$. Moreover, such a basis can be computed in time polynomial in $\|\lambda_1\| + \dots + \|\lambda_m\|$.*

The First-Order Theory of the Reals A sentence in the first-order theory of the reals is of the form $Q_1x_1 \cdots Q_mx_m\varphi(x_1, \dots, x_m)$ where each Q_i is a quantifier (\exists or \forall), each x_i is a real valued variable, and φ is a boolean combination of atomic predicates of the form $p(x_1, \dots, x_m) \sim 0$ for some $p \in \mathbb{Z}[x_1, \dots, x_m]$ and $\sim \in \{>, =\}$. The first-order theory of the reals admits quantifier elimination, a famous result due to Tarski [22], whose procedure unfortunately has non-elementary complexity. In this paper we consider only the case where the number of variables is uniformly bounded. Then, we can invoke the following result due to Renegar [20].

► **Theorem 10 (Renegar).** *Let $M \in \mathbb{N}$ be fixed. Let $\tau(\mathbf{y})$ be a formula of the first order theory of the reals. Assume that the number of (free and bound) variables in $\tau(\mathbf{y})$ is bounded by M . Denote the degree of $\tau(\mathbf{y})$ by d and the number of atomic predicates in $\tau(\mathbf{y})$ by n .*

There is a polynomial time (polynomial in $\|\tau(\mathbf{y})\|$) procedure which computes an equivalent quantifier-free formula

$$\chi(\mathbf{y}) = \bigvee_{i=1}^I \bigwedge_{j=1}^{J_i} h_{i,j}(\mathbf{y}) \sim_{i,j} 0$$

where each $\sim_{i,j}$ is either $>$ or $=$, with the following properties:

1. Each of I and J_i (for $1 \leq i \leq I$) is bounded by $(n+d)^{O(1)}$.
2. The degree of $\chi(\mathbf{y})$ is bounded by $(n+d)^{O(1)}$.
3. The height of $\chi(\mathbf{y})$ is bounded by $2^{\|\tau(\mathbf{y})\|(n+d)^{O(1)}}$.

Asymptotic Analysis We conclude this section with the following simple lemma from [17].

► **Proposition 11.** *Let $a \geq 2$ and $\epsilon \in (0, 1)$ be real numbers. Let $B \in \mathbb{Z}[x]$ have degree at most a^{D_1} and height at most $2^{a^{D_2}}$, and assume that $1/\epsilon \leq 2^{a^{D_3}}$ for some $D_1, D_2, D_3 \in \mathbb{N}$. Then there is $D_4 \in \mathbb{N}$ depending only on D_1, D_2, D_3 such that for all $n \geq 2^{a^{D_4}}$, $\frac{1}{B(n)} > (1 - \epsilon)^n$.*

3 Divergence

Recall from Theorem 1 that an LRS \mathbf{u} with dominant modulus ρ necessarily diverges in absolute value if $\rho > 1$. More precisely, if $\rho > 1$ then given $\epsilon > 0$ there exists a threshold N such that $|u_n| > \rho^{(1-\epsilon)n}$ for all $n > N$. However this result is *ineffective*—it is not known how to compute N given \mathbf{u} and ϵ .

In this section we derive *effective* divergence bounds for sequences that diverge to ∞ (i.e., sequences that both diverge in absolute value and that are ultimately positive). The bounds on divergence have the following form: for a divergent sequence \mathbf{u} with dominant modulus $\rho = 1$ we aim to show that for every $n > N$, $u_n > an^d$ for effective constants $a > 0, d \in \mathbb{N}$, and $N \in \mathbb{N}$. In case of a dominant modulus $\rho > 1$ we aim to show that for every $n > N$, $u_n > \frac{a\rho^n}{n^d}$ for effective constants $a > 0, d \in \mathbb{N}$, and $N \in \mathbb{N}$. Henceforth we refer to bounds of these respective forms as *divergence bounds*.⁵

In Section 3.1, we show how to compute effective divergence bounds of LRS up to certain orders. Then, in Section 3.2, we provide hardness results for the decidability of divergence.

⁵ Note that not only do we seek effective divergence bounds, but also that these bounds are asymptotically tighter than the bounds from Theorem 1 since for any fixed $d > 0$, it is clear that $a\rho^n/n^d$ eventually dominates $\rho^{(1-\epsilon)n}$ for any $\epsilon > 0$.

3.1 Effective Divergence is Solvable

In this section we prove the following theorems:

► **Theorem 12.** *There is a polynomial-time procedure that given a rational LRS of order at most 5 decides whether it diverges and, in case of divergence, outputs divergence bounds.*

► **Theorem 13.** *There is a polynomial-time procedure that, given a simple rational LRS of order at most 8, decides whether it diverges and, in case of divergence, outputs divergence bounds.*

The proofs of Theorems 12 and 13 build on techniques developed in [17, 16, 18], using a fine-grained analysis in the results thereof, along with some new ideas. To avoid unnecessary repetition, we sketch the main ideas of the proofs simultaneously.

Consider an LRS \mathbf{u} of order k . For uniformity, if \mathbf{u} is inhomogeneous, we homogenize it as per Proposition 3. Thus, either $k \leq 6$ or \mathbf{u} is simple and $k \leq 9$, where if $k = 6$ or if \mathbf{u} is simple and $k = 9$, then \mathbf{u} has a special structure according to Property 5.

As mentioned in Section 1, we can assume without loss of generality that \mathbf{u} is non-degenerate. Let ρ be the dominant modulus of \mathbf{u} , we also note that if $\rho < 1$, then $|u_n| \rightarrow 0$ as $n \rightarrow \infty$, and in particular the sequence does not diverge. Thus, we may assume $\rho \geq 1$. In addition, by Lemma 6, if \mathbf{u} does not have a real positive dominant root, then $u_n \not\rightarrow \infty$. Thus, we may assume a real positive dominant characteristic root ρ . Note that all other dominant roots must be complex, and come in conjugate pairs, since if $-\rho$ were a root, then \mathbf{u} would be degenerate.

Writing u_n as an exponential polynomial and dividing by ρ^n , we have

$$\frac{u_n}{\rho^n} = A(n) + \sum_{i=1}^m \left(C_i(n)\lambda_i^n + \overline{C_i(n)}\overline{\lambda_i}^n \right) + r(n) \quad (2)$$

where A is a real polynomial, C_i are non-zero complex polynomials, $\rho\lambda_i$ and $\rho\overline{\lambda_i}$ are conjugate pairs of non-real dominant characteristic roots of \mathbf{u} (so $|\lambda_i| = 1$), and $r(n)$ is an exponentially decaying function (possibly identically zero). More precisely, the degree of each of $A(n), C_1(n), \dots, C_m(n)$ is strictly smaller than the multiplicity of the corresponding characteristic root. We can assume that either $A(n) \not\equiv 0$ or $m \neq 0$. Indeed, otherwise we can consider the LRS $\langle \rho^n r(n) \rangle_{n=1}^{\infty}$, which is of lower order than \mathbf{u} .

In the following, if $A(n)$ (resp. $C_i(n)$ for some $1 \leq i \leq m$) is a constant, we denote it by A (resp. C_i).

We proceed to decide divergence by a case analysis of Equation (2).

Case 1: $\rho = 1$ and $A(n) = A$ is a constant

Note that in this case, $\frac{u_n}{\rho^n} = u_n$. Since A is a constant, then it does not affect the divergence of \mathbf{u} . We claim that $u_n \not\rightarrow \infty$. Indeed, by Lemma 6, the expression $\sum_{i=1}^m \left(C_i(n)\lambda_i^n + \overline{C_i(n)}\overline{\lambda_i}^n \right)$ becomes negative infinitely often (regardless of whether $C_i(n)$ are constants or polynomials), whereas the effect of $r(n)$ is exponentially decreasing. Thus, \mathbf{u} does not diverge.

Case 2: $\rho = 1$, $A(n)$ is not a constant, and every C_i is a constant

In this case we can rewrite Equation (2) as

$$u_n = A(n) + \sum_{i=1}^m \left(C_i\lambda_i^n + \overline{C_i}\overline{\lambda_i}^n \right) + r(n) \quad (3)$$

Since $|\lambda_i| = 1$ for all i , and since $r(n)$ is exponentially decreasing, then clearly $u_n \rightarrow \infty$ iff the leading coefficient of $A(n)$ is positive.

Recall that since $\rho = 1$, then if \mathbf{u} diverges, there exist $N, d \in \mathbb{N}$ and $a > 0$ such that $u_n \geq an^d$ for all $n > N$. We now show how to effectively compute N, d , and a .

From Proposition 7, we can compute in polynomial time $\epsilon \in (0, 1)$ and $N_1 \in \mathbb{N}$ such that $r(n) < (1 - \epsilon)^n < 1$ for all $n > N_1$. We thus have that $u_n \geq A(n) - 2 \sum_{i=1}^m |C_i| - 1$, and we can easily compute $N_2 \in \mathbb{N}$ and $a \in \mathbb{Q}$ (depending on the coefficients of $A(n)$) such that for all $n > N_2$ we have $A(n) - 2 \sum_{i=1}^m |C_i| - 1 \geq an^d$, where d is the degree of $A(n)$. Taking $N = \max\{N_1, N_2\}$, we conclude this case.

Case 3: $\rho = 1$, $A(n)$ is not a constant, and there exists a non-constant $C_i(n)$

We notice that if there exists a non-constant $C_i(n)$, it follows by Property 5 that \mathbf{u} is not obtained by homogenizing a simple LRS. That is, we are in the case where $k \leq 6$. In the notations of Equation (2), we then have that $m = 1$, $A(n)$ is linear, $C_1(n)$ is linear, and $r(n) \equiv 0$. Indeed, this corresponds to the case where the characteristic roots of u_n are $1, \lambda, \bar{\lambda}$, each with multiplicity 2. Let $A(n) = a_1n + b_1$ and $C_1(n) = a_2n + b_2$, then we can write

$$u_n = a_1n + b_1 + (a_2n + b_2)\lambda^n + (\bar{a}_2n + \bar{b}_2)\bar{\lambda}^n = n(a_1 + a_2\lambda^n + \bar{a}_2\bar{\lambda}^n) + (b_1 + b_2\lambda^n + \bar{b}_2\bar{\lambda}^n)$$

Since $|(b_1 + b_2\lambda^n + \bar{b}_2\bar{\lambda}^n)|$ is bounded, then u_n diverges iff $n(a_1 + a_2\lambda^n + \bar{a}_2\bar{\lambda}^n)$ diverges. Let $\theta = \arg \lambda$ and $\varphi = \arg a_2$. We have $n(a_1 + a_2\lambda^n + \bar{a}_2\bar{\lambda}^n) = n(a_1 + 2|a_2| \cos(n\theta + \varphi))$.

Observe that since \mathbf{u} is non-degenerate, then θ is not a rational multiple of π . It follows that $\{[n\theta + \varphi]_{2\pi} : n \in \mathbb{N}\}$ (where $[x]_{2\pi} = x - 2\pi j$ where j is the unique integer such that $0 \leq x - 2\pi j < 2\pi$) is dense in $[0, 2\pi)$, so $\{\cos(n\theta + \varphi) : n \in \mathbb{N}\}$ is dense in $[-1, 1]$. Again, we split into cases.

- If $a_1 > 2|a_2|$, we have that u_n diverges. Then, we can compute in polynomial time a rational $\epsilon > 0$ and $N \in \mathbb{N}$ such that $a_1 - 2|a_2| > \epsilon$ and $n(a_1 + 2|a_2|) - (b_1 - 2|b_2|) > \epsilon n$ for all $n > N$. We then have that $u_n > \epsilon n$ for all $n > N$, thus concluding effective decidability of divergence in this case.

- If $a_1 < 2|a_2|$, then u_n does not diverge, as it becomes negative infinitely often, by the density argument above.

- The remaining case is when $a_1 = 2|a_2|$, and the expression above becomes $na_1(1 + \cos(n\theta + \varphi))$. We show that in this case, u_n does not diverge.

By Taylor approximation, for every $x \in (-\pi, \pi]$ it holds that $1 - \cos(x) \leq \frac{x^2}{2}$. For $n \in \mathbb{N}$, write $\Lambda(n) = n\theta + \varphi - (2j + 1)\pi$, where $j \in \mathbb{Z}$ is the unique integer such that $-\pi < \Lambda(n) \leq \pi$. We now have that

$$na_1(1 + \cos(n\theta + \varphi)) = na_1(1 - \cos(n\theta + \varphi + \pi)) = na_1(1 - \cos(\Lambda(n))) < na_1 \frac{\Lambda(n)^2}{2}.$$

By Dirichlet's Approximation Theorem, we have that $|\Lambda(n)| < \frac{t}{n}$ for infinitely many values of n , where t is a constant depending on φ . Thus, we have $na_1 \frac{\Lambda(n)^2}{2} < \frac{a_1 t^2}{2n}$ for infinitely many values of n . It follows that u_n is infinitely often bounded by a constant, so it does not diverge.

Case 4: $\rho > 1$ and there exists a non-constant $C_i(n)$

As in Case 3, it holds that $k \leq 6$. Moreover, since $\rho > 1$, then whether or not \mathbf{u} was obtained by homogenization, the characteristic root 1 (if it exists) is captured in $r(n)$. Therefore, we have that $m = 1$, C_1 is linear, and $A(n) = A$ is constant. Let C_1 have leading coefficient

$b \neq 0$. By Lemma 6, there exists $\epsilon > 0$ such that $b\lambda^n + \overline{b\lambda}^n < -\epsilon$ infinitely often. Then $C_1(n)\lambda_1^n + \overline{C_1(n)\lambda_1}^n$ (and hence u_n) is unbounded below, so u_n does not diverge.

Case 5: $\rho > 1$, $A(n)$ is not a constant, and every C_i is a constant

Since $A(n)$ is not a constant and $\rho > 1$, this case may only arise for $k \leq 6$ and $m \leq 1$. We write

$$\frac{u_n}{\rho^n} = A(n) + C_1\lambda_1^n + \overline{C_1\lambda_1}^n + r(n).$$

where if $m = 0$ then take $C_1 = 0$.

If $A(n)$ has a negative leading coefficient, then u_n is unbounded from below, and in particular u_n does not diverge.

If $A(n)$ has a positive leading coefficient, we can compute in polynomial time $N_0 \in \mathbb{N}$ and a rational $\epsilon_0 > 0$ such that $A(n) - 2|C_1| > 2\epsilon_0$ for all $n > N_0$. By Proposition 7, we can also compute in polynomial time $N_1 \in \mathbb{N}$ and $\epsilon_1 \in (0, 1)$ such that $|r(n)| < (1 - \epsilon_1)^n$ for all $n > N_1$. Taking $N_2 \geq \log_{1-\epsilon_1} \epsilon_0$, we have that for all $n > \max\{N_0, N_1, N_2\}$, $|r(n)| < \epsilon_0$, and thus

$$\frac{u_n}{\rho^n} \geq A(n) - 2|C_1| + r(n) \geq A(n) - 2|C_1| - \epsilon_0 > 2\epsilon_0 - \epsilon_0 = \epsilon_0.$$

Thus we have $u_n \geq \epsilon_0\rho^n$ for all $n > \max\{N_0, N_1, N_2\}$, which immediately yields effective divergence bounds in this case.

Case 6: $\rho > 1$, $A(n) = A$ is a constant, and every C_i is a constant

This case is the most involved, and utilizes deep mathematical results. Our proof works along the lines of [18]. For completeness, the full proof can be found in Appendix A.

We rewrite Equation (2) as

$$\frac{u_n}{\rho^n} = A + \sum_{i=1}^m \left(C_i\lambda_i^n + \overline{C_i\lambda_i}^n \right) + r(n) \quad (4)$$

Observe that $m \leq 3$. Indeed, if $k \leq 8$ this is trivial, and if $k = 9$ then by Property 5, 1 must be a non-dominant characteristic root of \mathbf{u} , so $r(n) \neq 0$ and thus $m \leq 3$.

In the following, we handle the case $m = 3$. The cases where $m < 3$ are similar and slightly simpler.

Let $L = \{(v_1, \dots, v_3) \in \mathbb{Z}^3 : \lambda^{v_1} \dots \lambda^{v_3} = 1\}$, and let $\{\ell_1, \dots, \ell_p\}$ be a basis for L of cardinality p . Write $\ell_q = (\ell_{q,1}, \dots, \ell_{q,3})$ for $1 \leq q \leq p$. From Theorem 9, such a basis can be computed in polynomial time, and moreover – each $\ell_{q,j}$ may be assumed to have magnitude polynomial in $\|u\|$.

Consider the set $\mathbb{T} = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1| = |z_2| = |z_3| = 1 \text{ and for each } 1 \leq q \leq p, z_1^{\ell_{q,1}} z_2^{\ell_{q,2}} z_3^{\ell_{q,3}} = 1\}$.

Define $h : \mathbb{T} \rightarrow \mathbb{R}$ by setting $h(z_1, z_2, z_3) = \sum_{i=1}^3 (C_i z_i + \overline{C_i z_i})$, so that for every $n \in \mathbb{N}$, $\frac{u_n}{\rho^n} = A + h(\lambda_1^n, \lambda_2^n, \lambda_3^n) + r(n)$. By Kronecker's theorem on inhomogeneous Diophantine approximation [7], the set $\{\lambda_1^n, \lambda_2^n, \lambda_3^n : n \in \mathbb{N}\}$ is a dense subset of \mathbb{T} . Since h is continuous, it follows that $\inf \{h(\lambda_1^n, \lambda_2^n, \lambda_3^n) : n \in \mathbb{N}\} = \min h|_{\mathbb{T}} = \mu$ for some $\mu \in \mathbb{R}$.

In the full proof, we show that μ is algebraic, computable in polynomial time, with $\|\mu\| = \|\mathbf{u}\|^{O(1)}$.

We now split to cases according to the sign of $A + \mu$.

- If $A + \mu < 0$, then \mathbf{u} is infinitely often negative, and does not diverge.

42:10 Effective Divergence Analysis for Linear Recurrence Sequences

- If $A + \mu > 0$, then \mathbf{u} diverges, and we obtain an effective bound similarly to Case 5.
- It remains to analyze the case where $A + \mu = 0$. To this end, let $\lambda_j = e^{i\theta_j}$ and $C_j = |C_j|e^{i\varphi_j}$ for $1 \leq j \leq 3$. From Equation (4) we have

$$\frac{u_n}{\rho^n} = A + \sum_{j=1}^3 2|C_j| \cos(n\theta_j + \varphi_j) + r(n)$$

We further assume that all the C_j are non-zero. Indeed, if this does not hold, we can recast our analysis in lower dimension.

In the full proof, we use zero-dimensionality results to show that h achieves its minimum μ over \mathbb{T} only at a finite set of points $Z = \{(\zeta_1, \zeta_2, \zeta_3) \in \mathbb{T} : h(\zeta_1, \zeta_2, \zeta_3) = \mu\}$.

We concentrate on the set Z_1 of first coordinates of Z . Write

$$\tau_1(x) = \exists z_1 (\text{Re}(z_1) = x \wedge z_1 \in Z_1)$$

$$\tau_2(y) = \exists z_1 (\text{Im}(z_1) = y \wedge z_1 \in Z_1)$$

By rewriting these formulas in the first order theory of the reals, we are able to show, using Theorem 10, that any $\zeta_1 = \hat{x} + i\hat{y} \in Z_1$ is algebraic, and moreover satisfies $\|\zeta_1\| = \|\mathbf{u}\|^{O(1)}$. In addition, we show that the cardinality of Z_1 is at most polynomial in $\|\mathbf{u}\|$.

Since λ_1 is not a root of unity, for each $\zeta_1 \in Z_1$ there is at most one value of n such that $\lambda_1^n = \zeta_1$. Theorem 9 then entails that this value (if it exists) is at most $M = \|\mathbf{u}\|^{O(1)}$, which we can take to be uniform across all $\zeta_1 \in Z_1$. We can now invoke Corollary 8 to conclude that, for $n > M$, and for all $\zeta_1 \in Z_1$, we have

$$|\lambda_1^n - \zeta_1| > \frac{1}{n\|\mathbf{u}\|^D} \tag{5}$$

where $D \in \mathbb{N}$ is some absolute constant.

Let $b > 0$ be minimal such that the set

$$\left\{ z_1 \in \mathbb{C} : |z_1| = 1 \text{ and, for all } \zeta_1 \in Z_1, |z_1 - \zeta_1| \geq \frac{1}{b} \right\}$$

is non empty. Thanks to our bounds on the cardinality of Z_1 , we can use the first-order theory of the reals, together with Theorem 10, to conclude that b is algebraic and $\|b\| = \|\mathbf{u}\|^{O(1)}$.

Define the function $g : [b, \infty) \rightarrow \mathbb{R}$ as follows:

$$g(x) = \min\{h(z_1, z_2, z_3) - \mu : (z_1, z_2, z_3) \in \mathbb{T} \text{ and for all } \zeta_1 \in Z_1, |z_1 - \zeta_1| \geq \frac{1}{x}\}.$$

In the full proof we show that we can compute in polynomial time a polynomial $P \in \mathbb{Z}[x]$ such that, for all $x \in [b, \infty)$,

$$g(x) \geq \frac{1}{P(x)} \tag{6}$$

with $\|P\| = \|\mathbf{u}\|^{O(1)}$

By Proposition 7 we can find $\epsilon \in (0, 1)$ and $N = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for all $n > N$, we have $|r(n)| < (1 - \epsilon)^n$, and moreover $1/\epsilon = 2^{\|\mathbf{u}\|^{O(1)}}$. In addition, by Proposition 11, there is $N' = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for every $n \geq N'$

$$\frac{1}{2P(n\|\mathbf{u}\|^D)} > (1 - \epsilon)^n. \tag{7}$$

Combining Equations (4)–(7), we get

$$\begin{aligned} \frac{u_n}{\rho^n} &= A + h(\lambda_1^n, \lambda_2^n, \lambda_3^n) + r(n) \geq -\mu + h(\lambda_1^n, \lambda_2^n, \lambda_3^n) - (1 - \epsilon)^n \geq g(n^{\|\mathbf{u}\|^D}) - (1 - \epsilon)^n \\ &\geq \frac{1}{P(n^{\|\mathbf{u}\|^D})} - (1 - \epsilon)^n = \frac{1}{2P(n^{\|\mathbf{u}\|^D})} + \frac{1}{2P(n^{\|\mathbf{u}\|^D})} - (1 - \epsilon)^n \geq \frac{1}{2P(n^{\|\mathbf{u}\|^D})} \end{aligned}$$

provided $n > \max\{M, N, N'\}$. We thus have that $\frac{u_n}{\rho^n}$ is eventually lower bounded by an inverse polynomial and hence we have effective divergence bounds in this case.

Finally, Cases 1–6 allow us to conclude both Theorem 12 and Theorem 13.

3.2 Hardness of Divergence

We now turn to show lower bounds for the divergence problem. Surprisingly, our lower bounds hold already for homogeneous LRS, and for the divergence decision problem, even without requiring effectively computable bounds.

In [16], it is shown that the Ultimate Positivity problem for homogeneous LRS of order at least 6 is hard, in the sense that if Ultimate Positivity is decidable for such LRS, then certain hard open problems in Diophantine approximation become solvable. We show hardness of divergence for homogeneous LRS of order at least 6 by reducing from Ultimate Positivity.

► **Theorem 14.** *Ultimate Positivity is reducible to Divergence.*

Proof. We show a reduction from the Ultimate Positivity problem for non-degenerate LRS of order 6, shown to be hard in [16]. The key ingredient in the reduction is Theorem 1.

Consider a non-degenerate homogeneous LRS $\langle u_n \rangle$ of order 6 with dominant modulus ρ , and let $\mu = \max\left\{2, \frac{2}{\rho}\right\}$, then the sequence $v_n = \mu^n u_n$ is a non-degenerate homogeneous LRS of order 6 with dominant modulus $\mu\rho \geq 2$. By Theorem 1, taking $\epsilon = \frac{1}{2}$, it follows that there exists $N \in \mathbb{N}$ such $|v_n| \geq 2^{n/2}$ for every $n > N$. It immediately follows that v_n is ultimately positive iff $v_n \rightarrow \infty$. Clearly, however, v_n and u_n have the same sign, and therefore u_n is ultimately positive iff v_n diverges, and we are done. ◀

4 Positivity and Ultimate Positivity

In this section we study the Positivity and Ultimate Positivity problems for inhomogeneous LRS. These problems were studied in [16, 17, 18] for homogeneous LRS. Using Proposition 3 and some careful analysis, we extend the decidability results to the inhomogeneous case.

We start by citing some results from [16, 17, 18], split to upper and lower bounds.

► **Theorem 15** (Upper Bounds from [16, 17, 18]).

1. *Positivity and Ultimate Positivity are decidable for homogeneous LRS of order 5 or less with complexities in $\mathbf{coNP}^{\mathbf{PosSLP}}$ and \mathbf{PTIME} , respectively.*
2. *Positivity is decidable for simple homogeneous LRS of order 9 or less with complexity in $\mathbf{coNP}^{\mathbf{PosSLP}}$.*
3. *Ultimate Positivity is decidable for simple homogeneous LRS of any order with complexity in \mathbf{PTIME} .*
4. *Effective Ultimate Positivity is solvable for simple homogeneous LRS of order 9 or less with complexity in \mathbf{PTIME} .*

The following notion of hardness will be made precise in Section 4.2.

► **Theorem 16** (Lower Bounds from [16, 17, 18]). *Positivity and Ultimate Positivity for LRS of order at least 6 are hard with respect to certain hard open problems in Diophantine approximation.*

4.1 Upper Bounds

We proceed to prove analogous results to Theorem 15 for inhomogeneous LRS.

Theorem 15(1.) along with Proposition 3 readily give us the following:

► **Theorem 17.** *Positivity and Ultimate Positivity are decidable for inhomogeneous LRS of order 4 or less, with complexity in $\text{coNP}^{\text{PosSLP}}$.*

For simple LRS, things become more involved, as Proposition 3 does not preserve simplicity. However, Property 5 shows that simplicity is almost preserved, up to the multiplicity of the characteristic root 1. As we now show, this is sufficient to obtain upper bounds for inhomogeneous simple LRS.

We start by addressing effective Ultimate Positivity, which we then use for addressing Positivity.

► **Theorem 18.** *Effective Ultimate Positivity is solvable in polynomial time for simple inhomogeneous LRS of order 8 or less.*

Proof. Let \mathbf{v} be a simple, non-degenerate, inhomogeneous LRS of order 8 or less, and consider the homogeneous LRS $\mathbf{u} = \text{HOM}(\mathbf{v})$. By Proposition 3, \mathbf{u} is of order at most 9. If \mathbf{u} is a simple LRS, then by [17] we can effectively decide its Ultimate Positivity. We hence assume that \mathbf{u} is not simple.

By Property 5, it follows that the characteristic roots of \mathbf{u} all have multiplicity 1, apart from the characteristic root 1 which has multiplicity 2. Consider the dominant modulus ρ of \mathbf{u} . If $\rho > 1$, then by writing the exponential polynomial of \mathbf{u} , we have

$$\frac{u_n}{\rho^n} = a + \sum_{i=1}^m (c_i \lambda_i^n + \overline{c_i} \overline{\lambda_i^n}) + r(n) \tag{8}$$

with $a \in \mathbb{R}$, $c_i \in \mathbb{C} \setminus \mathbb{R}$ and $|\lambda_i| = 1$ for every $1 \leq i \leq m$, and $|r(n)|$ exponentially decaying. Crucially, since 1 is not a dominant characteristic root, its effect is enveloped in $r(n)$. Specifically, we observe that the analysis of effective Ultimate Positivity in [17] only relies on Proposition 7. Since this holds in the case at hand, we can effectively decide Ultimate Positivity when 1 is not a dominant characteristic root.

Finally, if 1 is a dominant characteristic root, the exponential polynomial of \mathbf{u} can be written as

$$u_n = A(n) + \sum_{i=1}^m (c_i \lambda_i^n + \overline{c_i} \overline{\lambda_i^n}) + r(n). \tag{9}$$

We observe that in this case, u_n is ultimately positive iff it diverges (indeed, clearly $|u_n| \rightarrow \infty$). Thus, we can reduce the problem to divergence, and proceed with the analysis as in Section 3 Case 2.

This concludes the proof that Ultimate Positivity is effectively decidable for simple inhomogeneous LRS of order at most 8. ◀

Similarly to Theorem 18, we are able to conclude the following result, whose proof can be found in Appendix B.1.

► **Theorem 19.** *Ultimate Positivity is decidable in polynomial time for simple inhomogeneous LRS of any order.*

Finally, using Theorem 18, we can solve the Positivity problem (see Appendix B.2 for the proof).

► **Theorem 20.** *Positivity is decidable for simple inhomogeneous LRS of order 8 or less, with complexity in $\text{coNP}^{\text{PosSLP}}$.*

4.2 Lower Bounds

We now turn to study lower bounds, proving analogous results to Theorem 16 for inhomogeneous LRS. Similarly to [17], the hardness results we present are with respect to long standing open problems in Diophantine approximation. Before stating our results, we require some definitions from Diophantine approximation. We refer the reader to [14, 17] for comprehensive references.

For any $x \in \mathbb{R}$, we define the *Lagrange constant* of x as

$$L_\infty(x) = \inf\{c \in \mathbb{R} : |x - \frac{n}{m}| \leq \frac{c}{m^2} \text{ for infinitely many } m, n \in \mathbb{Z}\}$$

and the *approximation type* of x as

$$L(x) = \inf\{c \in \mathbb{R} : |x - \frac{n}{m}| \leq \frac{c}{m^2} \text{ for some } m, n \in \mathbb{Z}\}$$

For the vast majority of transcendental numbers, the Lagrange constant and the approximation type are unknown, despite significant work [9, 17], and the problem of computing them is a major open problem. In the following, we show that the decidability of Ultimate Positivity (resp. Positivity) for inhomogeneous LRS of order 5 or more would imply a major breakthrough in computing the Lagrange constant (resp. approximation type) for a large class of transcendental numbers.

► **Theorem 21.** *If Ultimate Positivity is decidable for inhomogeneous rational LRS of order at least 5 then there is an algorithm that computes the Lagrange constant of any number $\theta/2\pi$ such that $e^{i\theta}$ has rational real and imaginary parts.*

Proof. In [17], it is shown that deciding Ultimate Positivity of the homogeneous LRS of order 6 given by

$$u_n = r \sin n\theta - n(1 - \cos n\theta) \text{ and } v_n = -r \sin n\theta - n(1 - \cos n\theta)$$

for every $r \in \mathbb{Q}$ such that $r > 0$ and $\theta \in (0, 2\pi)$ such that $e^{i\theta}$ has rational real and imaginary parts would allow one to compute $L_\infty(\theta/2\pi)$.

We observe that both sequences u_n and v_n fall under the premise of Proposition 4. Thus, by applying Proposition 4, we obtain an equivalent inhomogeneous LRS of order 5, concluding the proof. ◀

A similar proof, using the results of [17], gives us also the following theorem.

► **Theorem 22.** *If Positivity is decidable for inhomogeneous rational LRS of order at least 5 then there is an algorithm that computes the approximation type of any number $\theta/2\pi$ such that $e^{i\theta}$ has rational real and imaginary parts.*

References

- 1 S. Akshay, Timos Antonopoulos, Joël Ouaknine, and James Worrell. Reachability problems for Markov chains. *Inf. Process. Lett.*, 115(2):155–158, 2015.
- 2 Eric Allender, Peter Bürgisser, Johan Kjeldgaard-Pedersen, and Peter Bro Miltersen. On the complexity of numerical analysis. *SIAM J. Comput.*, 38(5):1987–2006, 2009.

- 3 Alan Baker and Gisbert Wüstholz. Logarithmic forms and group varieties. *J. reine angew. Math.*, 442(19-62):3, 1993.
- 4 W. J. Baumol. *Economic Dynamics. An Introduction*. Macmillan, 1970.
- 5 Jacek Bochnak, Michel Coste, and Marie-Françoise Roy. *Real algebraic geometry*, volume 36. Springer Science & Business Media, 2013.
- 6 Mark Braverman. Termination of integer linear programs. In *Computer Aided Verification, 18th International Conference, CAV 2006, Seattle, WA, USA, August 17-20, 2006, Proceedings*, pages 372–385, 2006.
- 7 John W.S. Cassels. *An Introduction to Diophantine Approximation*. Cambridge University Press, 1965.
- 8 Henri Cohen. *A course in computational algebraic number theory*, volume 138. Springer Science & Business Media, 2013.
- 9 Thomas W Cusick and Mary E Flahive. *The Markoff and Lagrange spectra*. Number 30. American Mathematical Soc., 1989.
- 10 Graham Everest, Alfred J. van der Poorten, Igor E. Shparlinski, and Thomas Ward. *Recurrence Sequences*, volume 104 of *Mathematical surveys and monographs*. American Mathematical Society, 2003.
- 11 J.-H. Evertse. On sums of S-units and linear recurrences. *Compositio Math.*, 53(2):225–244, 1984.
- 12 David W Masser. Linear relations on algebraic groups. *New Advances in Transcendence Theory*, pages 248–262, 1988.
- 13 M. Mignotte. A note on linear recursive sequences. *J. Austral. Math. Soc.*, 20(2):242–244, 1975.
- 14 Ivan Morton Niven. *Diophantine approximations*. Courier Corporation, 2008.
- 15 Joël Ouaknine and James Worrell. Effective positivity problems for simple linear recurrence sequences. *CoRR*, abs/1309.1550, 2013. URL: <http://arxiv.org/abs/1309.1550>.
- 16 Joël Ouaknine and James Worrell. On the positivity problem for simple linear recurrence sequences. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part II*, pages 318–329, 2014.
- 17 Joël Ouaknine and James Worrell. Positivity problems for low-order linear recurrence sequences. In *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014, Portland, Oregon, USA, January 5-7, 2014*, pages 366–379, 2014.
- 18 Joël Ouaknine and James Worrell. Ultimate positivity is decidable for simple linear recurrence sequences. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part II*, pages 330–341, 2014.
- 19 Joël Ouaknine and James Worrell. On linear recurrence sequences and loop termination. *SIGLOG News*, 2(2):4–13, 2015.
- 20 James Renegar. On the computational complexity and geometry of the first-order theory of the reals. part i: Introduction. preliminaries. the geometry of semi-algebraic sets. the decision problem for the existential theory of the reals. *Journal of symbolic computation*, 13(3):255–299, 1992.
- 21 T. N. Shorey and C. L. Stewart. On the Diophantine equation $ax^{2t} + bx^t y + cy^2 = d$ and pure powers in recurrence sequences. *Math. Scand.*, 52(1):24–36, 1983.
- 22 Alfred Tarski. A decision method for elementary algebra and geometry. *Bulletin of the American Mathematical Society*, 59, 1951.
- 23 A. J. van der Poorten and H. P. Schlickewei. *J. Austral. Math. Soc. Ser. A*, 51(1):154–170, 1991.

A Complete Proofs of Section 3

Before proceeding with the complete proofs of Theorems 12 and 13, we provide some additional mathematical tools,

A.1 Zero Dimensionality Results

The following zero-dimensionality lemmas are proved in [15].

► **Lemma 23.** *Let $a_1, \dots, a_m \in \mathbb{R}$ and $\varphi_1, \dots, \varphi_m \in \mathbb{R}$ be two collections of m real numbers, for $m \geq 1$, with each of the a_i non-zero, and let $l_1, \dots, l_m \in \mathbb{Z}$ be integers. Define $f, g : \mathbb{R}^m \rightarrow \mathbb{R}$ by setting $f(x_1, \dots, x_m) = \sum_{i=1}^m a_i \cos(x_i + \varphi_i)$ and $g(x_1, \dots, x_m) = \sum_{i=1}^m l_i x_i$. Assume that $g(x_1, \dots, x_m)$ is not of the form $l(x_i \pm x_j)$ for some non-zero $l \in \mathbb{Z}$ and indices $i \neq j$. Let $\psi \in \mathbb{R}$.*

Then the function f , subject to the constraint $g(x_1, \dots, x_m) = \psi$, achieves its minimum only finitely many times over the domain $[0, 2\pi)^m$.

► **Lemma 24.** *Let $\langle u_n \rangle$ be a non-degenerate simple LRS with dominant characteristic roots $\rho \in \mathbb{R}$ and $\gamma_1, \bar{\gamma}_1, \dots, \gamma_m, \bar{\gamma}_m \in \mathbb{C} \setminus \mathbb{R}$. Write $\lambda_i = \gamma_i/\rho$ for $1 \leq i \leq m$, and let $L = \{(v_1, \dots, v_m) \in \mathbb{Z}^m : \lambda_1^{v_1} \cdots \lambda_m^{v_m} = 1\}$. Let $\{\ell_1, \dots, \ell_{m-1}\}$ be a basis for L of cardinality $m - 1$, and write $\ell_j = (\ell_{j,1}, \dots, \ell_{j,m})$ for $1 \leq j \leq m - 1$. Let*

$$M = \begin{pmatrix} l_{1,1} & l_{1,2} & \cdots & l_{1,m-1} & l_{1,m} \\ l_{2,1} & l_{2,2} & \cdots & l_{2,m-1} & l_{2,m} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ l_{m-1,1} & l_{m-1,2} & \cdots & l_{m-1,m-1} & l_{m-1,m} \end{pmatrix}$$

Let $a_1, \dots, a_m \in \mathbb{R}$ and $\varphi_1, \dots, \varphi_m$ be two collections of m real numbers, with each of the a_i non-zero, and let $\mathbf{q} = (q_1, \dots, q_{m-1}) \in \mathbb{Z}^{m-1}$ be a column vector of $m - 1$ integers. Let us further write $\mathbf{x} = (x_1, \dots, x_m)$ to denote a column vector of m real-valued variables.

Then the function $f(x_1, \dots, x_m) = \sum_{i=1}^m a_i \cos(x_i + \varphi_i)$, subject to the constraint $M\mathbf{x} = 2\pi\mathbf{q}$, achieves its minimum at only finitely many points over the domain $[0, 2\pi)^m$.

A.2 Proof of Theorem 12

We initially prove Theorem 12 for homogeneous LRS. We then show how to handle the inhomogeneous case, using Property 5.

Consider an LRS $\mathbf{u} = \langle u_n \rangle_{n=1}^\infty$ of order $k \leq 5$ with dominant modulus ρ , and let $\epsilon > 0$ be a rational number. First, we note that without loss of generality, we can assume \mathbf{u} is non-degenerate, as we may decompose a degenerate sequence and recast analysis at lower orders. We also note that if $\rho < 1$, then $|u_n| \rightarrow 0$ as $n \rightarrow \infty$, and in particular the sequence does not diverge. Thus, we may assume $\rho \geq 1$.

By Lemma 6, if \mathbf{u} does not have a real positive dominant root, then $u_n \not\rightarrow \infty$. Thus, we may assume a real positive dominant root. Note that all other dominant roots must be complex, and come in conjugate pairs, since if $-\rho$ were a root, then \mathbf{u} would be degenerate.

Writing u_n as an exponential polynomial and dividing by ρ^n , we have

$$\frac{u_n}{\rho^n} = A(n) + \sum_{i=1}^m \left(C_i(n) \lambda_i^n + \overline{C_i(n)} \bar{\lambda}_i^n \right) + r(n) \quad (10)$$

where A is a real polynomial, C_i are non-zero complex polynomials, $\rho\lambda_i$ and $\rho\bar{\lambda}_i$ are conjugate pairs of non-real dominant roots of \mathbf{u} , and r is an exponentially decaying function (possibly

identically zero). We can assume that either $A \neq 0$ or $m \neq 0$. Indeed, otherwise we can consider the LRS $\langle \rho^n r(n) \rangle_{n=1}^\infty$, which is of lower order than \mathbf{u} .

We proceed to decide divergence by a case analysis of Equation (10).

Case 1: $\rho = 1$.

Note that in this case, $\frac{u_n}{\rho^n} = u_n$. If $A(n)$ is a constant, then it does not affect the divergence of \mathbf{u} . We claim that $u_n \not\rightarrow \infty$. Indeed, by Lemma 6, the expression $\sum_{i=1}^m (C_i(n)\lambda_i^n + \overline{C_i(n)}\overline{\lambda_i}^n)$ becomes negative infinitely often (regardless of whether C_i are constants or polynomials), whereas the effect of $r(n)$ is exponentially decreasing. Thus, \mathbf{u} does not diverge.

If $A(n)$ is not a constant, then $m \leq 1$. If $m = 0$, then clearly $u_n \rightarrow \infty$ iff the leading coefficient of $A(n)$ is positive. Otherwise, if $m = 1$, then C_1 is a constant, and thus $|C_1\lambda_1^n + \overline{C_1}\overline{\lambda_1}^n| \leq 2|C_1|$, and again $u_n \rightarrow \infty$ iff the leading coefficient of $A(n)$ is positive.

Recall that since $\rho = 1$, then if \mathbf{u} diverges, there exist $N, k \in \mathbb{N}$ such that $u_n \geq n^k$ for all $n > N$. We now show how to effectively compute N and k .

From Proposition 7, we can compute in polynomial time $\epsilon \in (0, 1)$ and $N_1 \in \mathbb{N}$ such that $r(n) < (1 - \epsilon)^n < 1$ for all $n > N_1$. We thus have that $u_n \geq A(n) - |C_1| - 1$, and we can easily compute $N_2 \in \mathbb{N}$ and $a \in \mathbb{Q}$ (depending on the coefficients of $A(n)$) such that for all $n > N_2$ we have $A(n) - |C_1| - 1 \geq an^k$, where k is the degree of $A(n)$, namely 1 or 2. Taking $N = \max\{N_1, N_2\}$, we conclude this case.

Case 2: $\rho > 1$ and there exists a non-constant C_i .

In this case, $m = 1$, C_1 is linear, and $A(n)$ is constant. Let C_1 have leading coefficient $b \neq 0$. By Lemma 6, there exists $\epsilon > 0$ such that $b\lambda^n + \overline{b}\overline{\lambda}^n < -\epsilon$ infinitely often. Then $C_1(n)\lambda_1^n + \overline{C_1(n)}\overline{\lambda_1}^n$ (and hence u_n) is unbounded below, so $u_n \not\rightarrow \infty$.

Case 3: $\rho > 1$ and every C_i is a nonzero constant.

In this case, $m \leq 2$. In the following, we set $m = 2$, as the cases where $m < 2$ are similar and slightly simpler.⁶

Let $L = \{(v_1, v_2) \in \mathbb{Z}^2 : \lambda^{v_1} \lambda^{v_2} = 1\}$, and let $\{\ell_1, \dots, \ell_p\}$ be a basis for L of cardinality p . Write $\ell_q = (\ell_{q,1}, \ell_{q,2})$ for $1 \leq q \leq p$. From Theorem 9, such a basis can be computed in polynomial time, and moreover – each $\ell_{q,j}$ may be assumed to have magnitude polynomial in $\|\mathbf{u}\|$.

Consider the set $\mathbb{T} = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| = |z_2| = 1 \text{ and for each } 1 \leq q \leq p, z_1^{\ell_{q,1}} z_2^{\ell_{q,2}} = 1\}$.

Define $h : \mathbb{T} \rightarrow \mathbb{R}$ by setting $h(z_1, z_2) = \sum_{i=1}^2 (C_i z_i + \overline{C_i} \overline{z_i})$, so that for every $n \in \mathbb{N}$, $\frac{u_n}{\rho^n} = A(n) + h(\lambda_1^n, \lambda_2^n) + r(n)$. Recall that the set $\{(\lambda_1^n, \lambda_2^n) : n \in \mathbb{N}\}$ is a dense subset of \mathbb{T} . Since h is continuous, it follows that $\inf\{h(\lambda_1^n, \lambda_2^n) : n \in \mathbb{N}\} = \min h|_{\mathbb{T}} = \mu$ for some $\mu \in \mathbb{R}$.

We now claim that μ is an algebraic number, computable in polynomial time, with $\|\mu\| = \|\mathbf{u}\|^{O(1)}$. We can represent μ via the following formula $\tau(y)$:

$$\exists(\zeta_1, \zeta_2) \in \mathbb{T} : [h(\zeta_1, \zeta_2) = y \wedge \forall(z_1, z_2) \in \mathbb{T}, y \leq h(z_1, z_2)].$$

Note that $\tau(y)$ is not a formula in the first-order theory of the reals, as it involves complex numbers. However, we can rewrite it as a sentence in the first-order theory of the reals

⁶ One may notice that taking $m = 2$ means that some of the cases we handle actually require order 6, e.g. when $A(n)$ is linear and $m = 2$. Still, the analysis covers all possible cases of order 5.

by representing the real and imaginary parts of each complex quantity and combine them using real arithmetic (see [16] for details). In addition, the obtained formula $\tau'(y)$ is of size polynomial in $\|\mathbf{u}\|$. By Theorem 10, we can then compute in polynomial time an equivalent quantifier-free formula

$$\chi(x) = \bigvee_{i=1}^I \bigwedge_{j=1}^{J_i} h_{i,j} \sim_{i,j} 0.$$

Recall that each $\sim_{i,j}$ is either $>$ or $=$. Now $\chi(x)$ must have a satisfiable disjunct, and since the satisfying assignment to y is unique (namely $y = \mu$), this disjunct must comprise at least one equality predicate. Since Theorem 10 guarantees that the degree and height of each $h_{i,j}$ are bounded by $\|\mathbf{u}\|^{O(1)}$ and $2^{\|\mathbf{u}\|^{O(1)}}$ respectively, we immediately conclude that μ is an algebraic number and with $\|\mu\| = \|\mathbf{u}\|^{O(1)}$.

We now split the analysis into several cases.

- If $A(n)$ is linear with negative leading coefficient, or if A is a constant and $A + \mu < 0$, then u_n is unbounded from below, and in particular $u_n \not\rightarrow \infty$.

- If $A(n)$ is linear with positive leading coefficient, or if A is a constant and $A + \mu > 0$, we can compute in polynomial time $N_0 \in \mathbb{N}$ and a rational $\epsilon_0 > 0$ such that $A(n) + \mu > 2\epsilon_0$ for all $n > N_0$. By Proposition 7, we can also compute in polynomial time $N_1 \in \mathbb{N}$ and $\epsilon_1 \in (0, 1)$ such that $|r(n)| < (1 - \epsilon_1)^n$ for all $n > N_1$. Taking $N_2 \geq \log_{1-\epsilon_1} \epsilon_0$, we have that for all $n > \max\{N_0, N_1, N_2\}$, $|r(n)| < \epsilon_0$, and thus

$$\frac{u_n}{\rho^n} = A(n) + h(\lambda_1, \dots, \lambda_m) + r(n) \geq A(n) + \mu - \epsilon_0 > 2\epsilon_0 - \epsilon_0 = \epsilon_0.$$

Thus we have $u_n \geq \epsilon_0 \rho^n$ for all $n > \max\{N_0, N_1, N_2\}$ and hence we have effective growth bounds in this case.

- If A is a constant and $A + \mu = 0$, things are more involved. Let $\lambda_j = e^{i\theta_j}$ and $C_j = |C_j|e^{i\varphi_j}$ for $1 \leq j \leq 2$. From Equation (10) we have

$$\frac{u_n}{\rho^n} = A + \sum_{j=1}^2 2|C_j| \cos(n\theta_j + \varphi_j) + r(n)$$

We further assume that all the C_j are non-zero. Indeed, if this does not hold, we can recast our analysis in lower dimension.

We now claim that h achieves its minimum μ only finitely many times over \mathbb{T} . To establish this claim, we proceed according to the cardinality p of the basis $\{\ell_1, \dots, \ell_p\}$ of L :

(i) We first consider the case in which $p = 1$, and handle the case $p = 0$ immediately afterwards. Let $\ell_1 = (\ell_{1,1}, \ell_{1,2}) \in \mathbb{Z}^2$ be the sole vector spanning L . For $x \in \mathbb{R}$, recall that we denote by $[x]_{2\pi}$ the distance from x to the closest integer multiple of 2π

Write

$$R = \{(x_1, x_2) \in [0, 2\pi)^2 : [\ell_{1,1}x_1 + \ell_{1,2}x_2]_{2\pi} = 0\}.$$

Clearly, for any $(x_1, x_2) \in [0, 2\pi)^2$, we have $(x_1, x_2) \in R$ iff $(e^{ix_1}, e^{ix_2}) \in \mathbb{T}$. Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by setting

$$f(x_1, x_2) = \sum_{j=1}^2 2|C_j| \cos(x_j + \varphi_j).$$

Clearly, for all $(x_1, x_2) \in [0, 2\pi)^2$ we have $f(x_1, x_2) = h(e^{ix_1}, e^{ix_2})$, and therefore the minimal of f over \mathbb{R} are in one-to-one correspondence with those of h over \mathbb{T} .

Define $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ by setting

$$g(x_1, x_2) = \ell_{1,1}x_1 + \ell_{1,2}x_2.$$

Note that $g(x_1, x_2)$ cannot be of the form $\ell(x_i - x_j)$, for nonzero $\ell \in \mathbb{Z}$ and $i \neq j$, otherwise $\lambda_i^\ell \lambda_j^{-\ell} = 1$, i.e., λ_i/λ_j would be a root of unity, contradicting the non-degeneracy of \mathbf{u} . Likewise, g cannot be of the form $\ell(x_i + x_j)$, otherwise $\lambda_i/\bar{\lambda}_j$ would be a root of unity.

Finally, observe that for $(x_1, x_2) \in [0, 2\pi)^2$, we have $(x_1, x_2) \in R$ iff $\ell_{1,1}x_1 + \ell_{1,2}x_2 = 2\pi q$ for some $q \in \mathbb{Z}$ with $|q| \leq |\ell_{1,1}| + |\ell_{1,2}|$. For each of these finitely many q , we can invoke Lemma 23 with f, g , and $\psi = 2\pi q$, to conclude that f achieves its minimum μ finitely many times over R , and therefore that h achieves the same minimum finitely many times over \mathbb{T} .

The case $p = 0$, i.e., in which there are no non-trivial integer multiplicative relationships among λ_1, λ_2 , is now a special case of the above, where we have $\ell_{1,1} = \ell_{1,2}$.

(ii) We observe that the case $p = 2$ cannot occur: indeed, a basis for L of dimension 2 would immediately entail that every λ_j is a root of unity.

This concludes the proof of the claim that h achieves its minimum at a finite number of points $Z = \{(\zeta_1, \zeta_2) \in \mathbb{T} : h(\zeta_1, \zeta_2) = \mu\}$.

We concentrate on the set Z_1 of first coordinates of Z . Write

$$\begin{aligned}\tau_1(x) &= \exists z_1 (\operatorname{Re}(z_1) = x \wedge z_1 \in Z_1) \\ \tau_2(y) &= \exists z_1 (\operatorname{Im}(z_1) = y \wedge z_1 \in Z_1)\end{aligned}$$

Similarly to our earlier construction, $\tau_1(x)$ is equivalent to a formula $t'_1(x)$ in the first-order theory of the reals, over a bounded number of real variables, with $\|\tau'_1(x)\| = \|\mathbf{u}\|^{O(1)}$. Thanks to Theorem 10, we then obtain an equivalent quantifier-free formula

$$\chi_1(x) = \bigvee_{i=1}^I \bigwedge_{j=1}^{J_i} h_{i,j} \sim_{i,j} 0.$$

Note that since there can only be finitely many $\hat{x} \in \mathbb{R}$ such that $\chi_1(\hat{x})$ holds, each disjunct of $\chi_1(\hat{x})$ must comprise at least one equality predicate, or can otherwise be entirely discarded as having no solution. A similar exercise can be carried out with $\tau_2(y)$ to obtain $\chi_2(y)$. The bounds on the degree and height of each $h_{i,j}$ in $\chi_1(x)$ and $\chi_2(y)$ then enables us to conclude that any $\zeta_1 = \hat{x} + i\hat{y} \in Z_1$ is algebraic, and moreover satisfies $\|\zeta_1\| = \|\mathbf{u}\|^{O(1)}$. In addition, bounds on I and J_i guarantee that the cardinality of Z_1 is at most polynomial in $\|\mathbf{u}\|$.

Since λ_1 is not a root of unity, for each $\zeta_1 \in Z_1$ there is at most one value of n such that $\lambda_1^n = \zeta_1$. Theorem 9 then entails that this value (if it exists) is at most $M = \|\mathbf{u}\|^{O(1)}$, which we can take to be uniform across all $\zeta_1 \in Z_1$. We can now invoke Corollary 8 to conclude that, for $n > M$, and for all $\zeta_1 \in Z_1$, we have

$$|\lambda_1^n - \zeta_1| > \frac{1}{n^{\|\mathbf{u}\|^D}} \tag{11}$$

where $D \in \mathbb{N}$ is some absolute constant.

Let $b > 0$ be minimal such that the set

$$\left\{ z_1 \in \mathbb{C} : |z_1| = 1 \text{ and, for all } \zeta_1 \in Z_1, |z_1 - \zeta_1| \geq \frac{1}{b} \right\}$$

is non empty. Thanks to our bounds on the cardinality of Z_1 , we can use the first-order theory of the reals, together with Theorem 10, to conclude that b is algebraic and $\|b\| = \|\mathbf{u}\|^{O(1)}$.

Define the function $g : [b, \infty) \rightarrow \mathbb{R}$ as follows:

$$g(x) = \min\{h(z_1, z_2) - \mu : (z_1, z_2) \in \mathbb{T} \text{ and for all } \zeta_1 \in Z_1, |z_1 - \zeta_1| \geq \frac{1}{x}\}.$$

It is clear that g is continuous and $g(x) > 0$ for all $x \in [b, \infty)$. Moreover, g can be translated in polynomial time into a function in the first-order theory of the reals over a bounded number of variables. It follows from Proposition 2.6.2 of [5] (invoked with the function $1/g$) that there is a polynomial $P \in \mathbb{Z}[x]$ such that, for all $x \in [b, \infty)$,

$$g(x) \geq \frac{1}{P(x)} \quad (12)$$

Moreover, an examination of the proof of [5, Prop. 2.6.2] reveals that P is obtained through a process which hinges on quantifier elimination. By Theorem 10, we are therefore able to conclude that $\|P\| = \|\mathbf{u}\|^{O(1)}$, a fact which relies among others on our upper bounds for $\|b\|$.

By Proposition 7 we can find $\epsilon \in (0, 1)$ and $N = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for all $n > N$, we have $|r(n)| < (1 - \epsilon)^n$, and moreover $1/\epsilon = 2^{\|\mathbf{u}\|^{O(1)}}$. In addition, by Proposition 11, there is $N' = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for every $n \geq N'$

$$\frac{1}{2P(n\|\mathbf{u}\|^D)} > (1 - \epsilon)^n. \quad (13)$$

Combining Equations (10)–(13), we get

$$\begin{aligned} \frac{u_n}{\rho^n} &= A + h(\lambda_1^n, \lambda_2^n) + r(n) \\ &\geq -\mu + h(\lambda_1^n, \lambda_2^n) - (1 - \epsilon)^n \\ &\geq g(n\|\mathbf{u}\|^D) - (1 - \epsilon)^n \\ &\geq \frac{1}{P(n\|\mathbf{u}\|^D)} - (1 - \epsilon)^n \\ &= \frac{1}{2P(n\|\mathbf{u}\|^D)} + \frac{1}{2P(n\|\mathbf{u}\|^D)} - (1 - \epsilon)^n \\ &\geq \frac{1}{2P(n\|\mathbf{u}\|^D)} \end{aligned}$$

provided $n > \max\{M, N, N'\}$. We thus have that $\frac{u_n}{\rho^n}$ is eventually lower bounded by an inverse polynomial and hence we have effective growth bounds in this case.

This concludes the decidability of divergence and computability of effective bounds on divergence for homogeneous LRS of order at most 5.

It remains to show how to handle inhomogeneous LRS of order at most⁷ 5. Consider an inhomogeneous LRS $\mathbf{v} = \langle v_n \rangle_{n=1}^\infty$ of order 5, and let $\mathbf{u} = \text{HOM}(\mathbf{v})$. Consider the dominant modulus ρ of u_n . If $\rho > 1$, then by property 5 the exponential polynomial of $\frac{u_n}{\rho^n}$ is the same as that in Equation (10). Thus, we can proceed with the case analysis of Case 2 and Case 3 without change. If $\rho = 1$, things become more involved. Consider the exponential polynomial

$$u_n = A(n) + \sum_{i=1}^m \left(C_i(n)(\lambda_i^n) + \overline{C_i(n)}(\overline{\lambda_i}^n) \right) + r(n) \quad (14)$$

where $|r(n)|$ is exponentially decaying and the λ_i are characteristic roots of modulus 1.

If $A(n)$ is constant, or if $A(n)$ is not a constant and all the C_i are constants (if there are any), then the same analysis of Case 1 applies here, *mutatis-mutandis*. Otherwise, the only possible case is where $A(n)$ is linear, $m = 1$, $C_1(n)$ is linear, and $r(n) \equiv 0$. Indeed, this

⁷ In fact, by property 5, LRS of order at most 4 can be handled by homogenization. Thus, it is enough to handle exactly order 5.

corresponds to the case where the characteristic roots of u_n are $1, \lambda, \bar{\lambda}$, each with multiplicity 2. Let $A(n) = a_1n + b_1$ and $C_1(n) = a_2n + b_2$, then we can write

$$u_n = a_1n + b_1 + (a_2n + b_2)\lambda^n + (\bar{a}_2n + \bar{b}_2)\bar{\lambda}^n = n(a_1 + a_2\lambda^n + \bar{a}_2\bar{\lambda}^n) + (b_1 + b_2\lambda^n + \bar{b}_2\bar{\lambda}^n)$$

Since $|(b_1 + b_2\lambda^n + \bar{b}_2\bar{\lambda}^n)|$ is bounded, then u_n diverges iff $n(a_1 + a_2\lambda^n + \bar{a}_2\bar{\lambda}^n)$ diverges. Let $\theta = \arg \lambda$ and $\varphi = \arg a_2$. We have $n(a_1 + a_2\lambda^n + \bar{a}_2\bar{\lambda}^n) = n(a_1 + 2|a_2| \cos(n\theta + \varphi))$.

Again, we split into cases.

- If $a_1 > 2|a_2|$, we have that u_n diverges. Then, we can compute in polynomial time a rational $\epsilon > 0$ and $N \in \mathbb{N}$ such that $a_1 - 2|a_2| > \epsilon$ and $n(a_1 + 2|a_2|) - (b_1 - 2|b_2|) > \epsilon n$ for all $n > N$. We then have that $u_n > \epsilon n$ for all $n > N$, thus concluding effective decidability of divergence in this case.

- If $a_1 < 2|a_2|$, then u_n does not diverge, as it becomes negative infinitely often.

- The remaining case is when $a_1 = 2|a_2|$, where the expression above becomes $na_1(1 + \cos(n\theta + \varphi))$. We show that in this case, u_n does not diverge. By Taylor approximation, for every $x \in (-\pi, \pi]$ it holds that $1 - \cos(x) \leq \frac{x^2}{2}$. For $n \in \mathbb{N}$, write $\Lambda(n) = n\theta + \varphi - (2j + 1)\pi$, where $j \in \mathbb{Z}$ is the unique integer such that $-\pi < \Lambda(n) \leq \pi$. We now have that

$$na_1(1 + \cos(n\theta + \varphi)) = na_1(1 - \cos(n\theta + \varphi + \pi)) = na_1(1 - \cos(\Lambda(n))) < na_1 \frac{\Lambda(n)^2}{2}$$

By Dirichlet's Approximation Theorem, we have that $|\Lambda(n)| < \frac{t}{n}$ for infinitely many values of n , where t is a constant depending on φ . Thus, we have $na_1 \frac{\Lambda(n)^2}{2} < \frac{a_1 t^2}{2n}$. It follows that u_n is infinitely often bounded by a constant, and does not diverge.

A.3 Proof of Theorem 13

As in the proof of Theorem 12, we start by considering the homogeneous case, and we let $\langle u_n \rangle$ be a non-degenerate simple LRS of order $d \leq 8$ with a real positive dominant characteristic root $\rho \geq 1$.

As before, we write

$$\frac{u_n}{\rho^n} = a + \sum_{i=1}^m (c_i \lambda_i^n + \bar{c}_i \bar{\lambda}_i^n) + r(n) \tag{15}$$

with $a \in \mathbb{R}$, $c_i \in \mathbb{C} \setminus \mathbb{R}$ for every $1 \leq i \leq m$, and $|r(n)|$ exponentially decaying. Note that since $d \leq 8$ and $a \in \mathbb{R}$, it follows that $0 \leq m \leq 3$. In the following, we consider the case where $m = 3$. The cases where $m < 3$ are very similar and slightly simpler, and are therefore omitted.

Observe that if $\rho = 1$, the sequence u_n is bounded, and therefore does not diverge. We hence assume $\rho > 1$.

Let $L = \{(v_1, \dots, v_m) \in \mathbb{Z}^m : \lambda^{v_1} \dots \lambda^{v_m} = 1\}$, and let $\{\ell_1, \dots, \ell_p\}$ be a basis for L of cardinality p . Write $\ell_{\mathbf{q}} = (\ell_{q,1}, \dots, \ell_{q,m})$ for $1 \leq q \leq p$. From Theorem 9, such a basis can be computed in polynomial time, and moreover – each $\ell_{q,j}$ may be assumed to have magnitude polynomial in $\|u\|$.

Consider the set $\mathbb{T} = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1| = |z_2| = |z_3| = 1 \text{ and for each } 1 \leq q \leq p, z_1^{\ell_{q,1}} z_2^{\ell_{q,2}} z_3^{\ell_{q,3}} = 1\}$.

Define $h : \mathbb{T} \rightarrow \mathbb{R}$ by setting $h(z_1, z_2, z_3) = \sum_{i=1}^3 (c_i z_i + \bar{c}_i \bar{z}_i)$, so that for every $n \in \mathbb{N}$, $\frac{u_n}{\rho^n} = a + h(\lambda_1^n, \lambda_2^n, \lambda_3^n) + r(n)$. Recall that the set $\{\lambda_1^n, \lambda_2^n, \lambda_3^n : n \in \mathbb{N}\}$ is a dense subset of \mathbb{T} . Since h is continuous, it follows that $\inf \{h(\lambda_1^n, \lambda_2^n, \lambda_3^n) : n \in \mathbb{N}\} = \min h|_{\mathbb{T}} = \mu$ for some $\mu \in \mathbb{R}$.

We now claim that μ is an algebraic number, computable in polynomial time, with $\|\mu\| = \|\mathbf{u}\|^{O(1)}$. We can represent μ via the following formula $\tau(y)$:

$$\exists(\zeta_1, \zeta_2, \zeta_3) \in \mathbb{T}[h(\zeta_1, \zeta_2, \zeta_3) = y \wedge \forall(z_1, z_2, z_3) \in \mathbb{T}, y \leq h(z_1, z_2, z_3)].$$

Note that $\tau(y)$ is not a formula in the first-order theory of the reals, as it involves complex numbers. However, we can rewrite it as a sentence in the first-order theory of the reals by representing the real and imaginary parts of each complex quantity and combine them using real arithmetic (see [16] for details). In addition, the obtained formula $\tau'(y)$ is of size polynomial in $\|\mathbf{u}\|$. By Theorem 10, we can then compute in polynomial time an equivalent quantifier-free formula

$$\chi(x) = \bigvee_{i=1}^I \bigwedge_{j=1}^{J_i} h_{i,j} \sim_{i,j} 0.$$

Recall that each $\sim_{i,j}$ is either $>$ or $=$. Now $\chi(x)$ must have a satisfiable disjunct, and since the satisfying assignment to y is unique (namely $y = \mu$), this disjunct must comprise at least one equality predicate. Since Theorem 10 guarantees that the degree and height of each $h_{i,j}$ are bounded by $\|\mathbf{u}\|^{O(1)}$ and $2\|\mathbf{u}\|^{O(1)}$ respectively, we immediately conclude that μ is an algebraic number and with $\|\mu\| = \|\mathbf{u}\|^{O(1)}$.

We now split to cases according to the sign of $a + \mu$.

- If $a + \mu < 0$, then \mathbf{u} is infinitely often negative, and does not diverge.
- If $a + \mu > 0$, then \mathbf{u} diverges, and it remains to show an effective bound. We can compute in polynomial time a rational $\epsilon_0 > 0$ such that $a + \mu > 2\epsilon_0$. By Proposition 7, we can also compute in polynomial time $N_1 \in \mathbb{N}$ and $\epsilon_1 \in (0, 1)$ such that $|r(n)| < (1 - \epsilon_1)^n$ for all $n > N_1$. Taking $N_2 \geq \log_{1-\epsilon_1} \epsilon_0$, we have that for all $n > \max\{N_1, N_2\}$, $|r(n)| < \epsilon_0$, and thus

$$\frac{u_n}{\rho^n} = A(n) + h(\lambda_1, \dots, \lambda_m) + r(n) \geq A(n) + \mu - \epsilon_0 > 2\epsilon_0 - \epsilon_0 = \epsilon_0$$

Thus $u_n > \epsilon_0 \rho^n$ for all $n > \max\{N_1, N_2\}$ and hence we have effective divergence bounds in this case.

- It remains to analyze the case where $a + \mu = 0$. To this end, let $\lambda_j = e^{i\theta_j}$ and $c_j = |c_j|e^{i\varphi_j}$ for $1 \leq j \leq 3$. From Equation (15) we have

$$\frac{u_n}{\rho^n} = a + \sum_{j=1}^3 2|c_j| \cos(n\theta_j + \varphi_j) + r(n)$$

We further assume that all the c_j are non-zero. Indeed, if this does not hold, we can recast our analysis in lower dimension.

We now claim that h achieves its minimum μ only finitely many times over \mathbb{T} . To establish this claim, we proceed according to the cardinality p of the basis $\{\ell_1, \dots, \ell_p\}$ of L :

- We first consider the case in which $p = 1$, and handle the case $p = 0$ immediately afterwards. Let $\ell_1 = (\ell_{1,1}, \ell_{1,2}, \ell_{1,3}) \in \mathbb{Z}^3$ be the sole vector spanning L . For $x \in \mathbb{R}$, recall that we denote by $[x]_{2\pi}$ the distance from x to the closest integer multiple of 2π . Write

$$R = \{(x_1, x_2, x_3) \in [0, 2\pi)^3 : [\ell_{1,1}x_1 + \ell_{1,2}x_2 + \ell_{1,3}x_3]_{2\pi} = 0\}.$$

Clearly, for any $(x_1, x_2, x_3) \in [0, 2\pi)^3$, we have $(x_1, x_2, x_3) \in R$ iff $(e^{ix_1}, e^{ix_2}, e^{ix_3}) \in \mathbb{T}$. Define $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ by setting

$$f(x_1, x_2, x_3) = \sum_{j=1}^3 2|c_j| \cos(x_j + \varphi_j).$$

Clearly, for all $(x_1, x_2, x_3) \in [0, 2\pi)^3$ we have $f(x_1, x_2, x_3) = h(e^{ix_1}, e^{ix_2}, e^{ix_3})$, and therefore the minima of f over \mathbb{R} are in one-to-one correspondence with those of h over \mathbb{T} .

Define $g : \mathbb{R}^3 \rightarrow \mathbb{R}$ by setting

$$g(x_1, x_2, x_3) = \ell_{1,1}x_1 + \ell_{1,2}x_2 + \ell_{1,3}x_3.$$

Note that $g(x_1, x_2, x_3)$ cannot be of the form $\ell(x_i - x_j)$, for nonzero $\ell \in \mathbb{Z}$ and $i \neq j$, otherwise $\lambda_i^\ell \lambda_j^{-\ell} = 1$, i.e., λ_i/λ_j would be a root of unity, contradicting the non-degeneracy of \mathbf{u} . Likewise, g cannot be of the form $\ell(x_i + x_j)$, otherwise $\lambda_i/\bar{\lambda}_j$ would be a root of unity.

Finally, observe that for $(x_1, x_2, x_3) \in [0, 2\pi)^3$, we have $(x_1, x_2, x_3) \in R$ iff $\ell_{1,1}x_1 + \ell_{1,2}x_2 + \ell_{1,3}x_3 = 2\pi q$ for some $q \in \mathbb{Z}$ with $|q| \leq |\ell_{1,1}| + |\ell_{1,2}| + |\ell_{1,3}|$. For each of these finitely many q , we can invoke Lemma 23 with f, g , and $\psi = 2\pi q$, to conclude that f achieves its minimum μ finitely many times over R , and therefore that h achieves the same minimum finitely many times over \mathbb{T} .

The case $p = 0$, i.e., in which there are no non-trivial integer multiplicative relationships among $\lambda_1, \lambda_2, \lambda_3$, is now a special case of the above, where we have $\ell_{1,1} = \ell_{1,2} = \ell_{1,3} = 0$.

(ii) We now turn to the case $p = 2$. We have $\ell_1 = (\ell_{1,1}, \ell_{1,2}, \ell_{1,3}) \in \mathbb{Z}^3$ and $\ell_2 = (\ell_{2,1}, \ell_{2,2}, \ell_{2,3}) \in \mathbb{Z}^3$ spanning L . Let \mathbf{x} denote the column vector (x_1, x_2, x_3) , and write

$$R = \{(x_1, x_2, x_3) \in [0, 2\pi)^3 : [\ell_1 \cdot \mathbf{x}]_{2\pi} = 0 \text{ and } [\ell_2 \cdot \mathbf{x}]_{2\pi} = 0\}.$$

Define $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ by setting $f(x_1, x_2, x_3) = \sum_{j=1}^3 2|c_j| \cos(x_j + \varphi_j)$. As before, the minima of f over R are in one-to-one correspondence with those of h over \mathbb{T} .

For $(x_1, x_2, x_3) \in [0, 2\pi)^3$, we have $[\ell_1 \cdot \mathbf{x}]_{2\pi} = 0$ and $[\ell_2 \cdot \mathbf{x}]_{2\pi} = 0$ iff there exist $q_1, q_2 \in \mathbb{Z}$, with $|q_1| \leq |\ell_{1,1}| + |\ell_{1,2}| + |\ell_{1,3}|$ and $|q_2| \leq |\ell_{2,1}| + |\ell_{2,2}| + |\ell_{2,3}|$ such that $\ell_1 \cdot \mathbf{x} = 2\pi q_1$ and $\ell_2 \cdot \mathbf{x} = 2\pi q_2$. For each of these finitely many $\mathbf{q} = (q_1, q_2)$, we can invoke Lemma 24 with $f, M = \begin{pmatrix} \ell_{1,1} & \ell_{1,2} & \ell_{1,3} \\ \ell_{2,1} & \ell_{2,2} & \ell_{2,3} \end{pmatrix}$, and \mathbf{q} , to conclude that f achieves its minimum μ finitely many times over R , and therefore that h achieves the same minimum finitely many times over \mathbb{T} .

(iii) Finally, we observe that the case $p = 3$ cannot occur: indeed, a basis for L of dimension 3 would immediately entail that every λ_j is a root of unity.

This concludes the proof of the claim that h achieves its minimum at a finite number of points $Z = \{(\zeta_1, \zeta_2, \zeta_3) \in \mathbb{T} : h(\zeta_1, \zeta_2, \zeta_3) = \mu\}$.

We concentrate on the set Z_1 of first coordinates of Z . Write

$$\tau_1(x) = \exists z_1 (\text{Re}(z_1) = x \wedge z_1 \in Z_1)$$

$$\tau_2(y) = \exists z_1 (\text{Im}(z_1) = y \wedge z_1 \in Z_1)$$

Similarly to our earlier constructions $\tau_1(x)$ is equivalent to a formula $t'_1(x)$ in the first-order theory of the reals, over a bounded number of real variables, with $\|\tau'_1(x)\| = \|\mathbf{u}\|^{O(1)}$. Thanks to Theorem 10, we then obtain an equivalent quantifier-free formula

$$\chi_1(x) = \bigvee_{i=1}^I \bigwedge_{j=1}^{J_i} h_{i,j} \sim_{i,j} 0.$$

Note that since there can only be finitely many $\hat{x} \in \mathbb{R}$ such that $\chi_1(\hat{x})$ holds, each disjunct of $\chi_1(\hat{x})$ must comprise at least one equality predicate, or can otherwise be entirely discarded as having no solution. A similar exercise can be carried out with $\tau_2(x)$. The bounds on the degree and height of each $h_{i,j}$ in $\chi_1(x)$ and $\chi_2(y)$ then enables us to conclude that any $\zeta_1 = \hat{x} + i\hat{y} \in Z_1$ is algebraic, and moreover satisfies $\|\zeta_1\| = \|\mathbf{u}\|^{O(1)}$. In addition, bounds on I and J_i guarantee that the cardinality of Z_1 is at most polynomial in $\|\mathbf{u}\|$.

Since λ_1 is not a root of unity, for each $\zeta_1 \in Z_1$ there is at most one value of n such that $\lambda_1^n = \zeta_1$. Theorem 9 then entails that this value (if it exists) is at most $M = \|\mathbf{u}\|^{O(1)}$, which we can take to be uniform across all $\zeta_1 \in Z_1$. We can now invoke Corollary 8 to conclude that, for $n > M$, and for all $\zeta_1 \in Z_1$, we have

$$|\lambda_1^n - \zeta_1| > \frac{1}{n\|\mathbf{u}\|^D} \quad (16)$$

where $D \in \mathbb{N}$ is some absolute constant.
Let $b > 0$ be minimal such that the set

$$\left\{ z_1 \in \mathbb{C} : |z_1| = 1 \text{ and, for all } \zeta_1 \in Z_1, |z_1 - \zeta_1| \geq \frac{1}{b} \right\}$$

is non empty. Thanks to our bounds on the cardinality of Z_1 , we can use the first-order theory of the reals, together with Theorem 10, to conclude that b is algebraic and $\|b\| = \|\mathbf{u}\|^{O(1)}$.

Define the function $g : [b, \infty) \rightarrow \mathbb{R}$ as follows:

$$g(x) = \min\{h(z_1, z_2, z_3) - \mu : (z_1, z_2, z_3) \in \mathbb{T} \text{ and for all } \zeta_1 \in Z_1, |z_1 - \zeta_1| \geq \frac{1}{x}\}.$$

It is clear that g is continuous and $g(x) > 0$ for all $x \in [b, \infty)$. Moreover, g can be translated in polynomial time into a function in the first-order theory of the reals over a bounded number of variables. It follows from Proposition 2.6.2 of [5] (invoked with the function $1/g$) that there is a polynomial $P \in \mathbb{Z}[x]$ such that, for all $x \in [b, \infty)$,

$$g(x) \geq \frac{1}{P(x)} \quad (17)$$

Moreover, an examination of the proof of [5, Prop. 2.6.2] reveals that P is obtained through a process which hinges on quantifier elimination. By Theorem 10, we are therefore able to conclude that $\|P\| = \|\mathbf{u}\|^{O(1)}$, a fact which relies among others on our upper bounds for $\|b\|$.

By Proposition 7 we can find $\epsilon \in (0, 1)$ and $N = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for all $n > N$, we have $|r(n)| < (1 - \epsilon)^n$, and moreover $1/\epsilon = 2^{\|\mathbf{u}\|^{O(1)}}$. In addition, by Proposition 11, there is $N' = 2^{\|\mathbf{u}\|^{O(1)}}$ such that for every $n \geq N'$

$$\frac{1}{2P(n\|\mathbf{u}\|^D)} > (1 - \epsilon)^n. \quad (18)$$

Combining Equations (15)–(18), we get

$$\begin{aligned} \frac{u_n}{\rho^n} &= a + h(\lambda_1^n, \lambda_2^n, \lambda_3^n) + r(n) \\ &\geq -\mu + h(\lambda_1^n, \lambda_2^n, \lambda_3^n) - (1 - \epsilon)^n \\ &\geq g(n\|\mathbf{u}\|^D) - (1 - \epsilon)^n \\ &\geq \frac{1}{P(n\|\mathbf{u}\|^D)} - (1 - \epsilon)^n \\ &= \frac{1}{2P(n\|\mathbf{u}\|^D)} + \frac{1}{2P(n\|\mathbf{u}\|^D)} - (1 - \epsilon)^n \\ &\geq \frac{1}{2P(n\|\mathbf{u}\|^D)} \end{aligned}$$

provided $n > \max\{M, N, N'\}$. We thus have that $\frac{u_n}{\rho^n}$ is eventually lower bounded by an inverse polynomial and hence we have effective divergence bounds in this case.

It remains to show how to handle inhomogeneous LRS of order at most 8. Consider an inhomogeneous LRS $\langle v_n \rangle$ of order at most 8, and let $u_n = \text{HOM}(v_n)$. Observe that by Property 5, u_n might not be a simple LRS. However, all its characteristic roots have multiplicity 1, apart from, possibly, the characteristic root 1.

Consider the dominant modulus ρ of u_n . If $\rho > 1$, then by property 5 the exponential polynomial of $\frac{u_n}{\rho^n}$ is of the same form as that in Equation (10), in the sense that $r(n)$ is still exponentially decaying. Thus, we can proceed with the analysis above without change. If $\rho = 1$, things become slightly more involved. Consider the exponential polynomial

$$u_n = A(n) + \sum_{i=1}^m (c_i \lambda_i^n + \overline{c_i} \overline{\lambda_i}^n) + r(n) \tag{19}$$

where $A(n)$ is either a constant or a polynomial, $c_i \in \mathbb{C} \setminus \mathbb{R}$ for every $1 \leq i \leq m$, $|r(n)|$ is exponentially decaying, and $0 \leq m \leq 4$. Observe that if $A(n)$ is a constant, then $|u_n|$ is bounded, and so does not diverge. In particular, if $m = 4$ then it has to be the case that $A(n)$ is constant. Thus, it suffices to consider the case where $m \leq 3$ and $A(n)$ is a polynomial.

In this case, similarly to Case 1 in the proof of Theorem 12, we have that \mathbf{u} diverges iff the leading coefficient of $A(n)$ is positive, and in this case the bounds are effectively computable.

This completes the proof of Theorem 13.

B Complete Proofs of Section 4

B.1 Proof of Theorem 19

Let \mathbf{v} be a simple, non-degenerate, inhomogeneous LRS, and consider the homogeneous LRS $\mathbf{u} = \text{HOM}(\mathbf{v})$. If \mathbf{u} is a simple LRS, then by [17] we can effectively decide its Ultimate Positivity. We assume henceforth that \mathbf{u} is not simple.

By Property 5, it follows that the characteristic roots of \mathbf{u} all have multiplicity 1, apart from the characteristic root 1 which has multiplicity 2. Consider the dominant modulus ρ of \mathbf{u} . If $\rho > 1$, then by writing the exponential polynomial of \mathbf{u} , we have

$$\frac{u_n}{\rho^n} = a + \sum_{i=1}^m (c_i \lambda_i^n + \overline{c_i} \overline{\lambda_i}^n) + r(n) \tag{20}$$

with $a \in \mathbb{R}$, $c_i \in \mathbb{C} \setminus \mathbb{R}$ and $|\lambda_i| = 1$ for every $1 \leq i \leq m$, and $|r(n)|$ exponentially decaying. Crucially, since 1 is not a dominant characteristic root, its effect is enveloped in $r(n)$. Specifically, we observe that the analysis of effective Ultimate Positivity in [18] only relies on Proposition 7. Since this holds in the case at hand, we can effectively decide Ultimate Positivity when 1 is not a dominant characteristic root.

Finally, if 1 is a dominant characteristic root, the exponential polynomial of \mathbf{u} can be written as

$$u_n = A(n) + \sum_{i=1}^m (c_i \lambda_i^n + \overline{c_i} \overline{\lambda_i}^n) + r(n) \tag{21}$$

We observe that in this case, u_n is ultimately positive iff it diverges (indeed, clearly $|u_n| \rightarrow \infty$). Thus, we can reduce the problem to divergence, and proceed with the analysis as in Section 3 Case 2. ◀

B.2 Proof of Theorem 20

Given the proof of Theorem 18, Positivity is now easily decidable: given an inhomogeneous simple LRS \mathbf{u} of order at most 8, decide if its ultimately positive, and if so - compute the bound from which it is ultimately positive. Then, deciding Positivity amounts to checking a finite number of elements.

Note that the bound computed in Theorem 18 is $N = 2^{\mathbf{u}^{O(1)}}$. This implies that checking whether an ultimately-positive LRS is *not* positive can be done using a *guess-and-check* procedure, and employing **PosSLP** in order to compute double exponential numbers. This yields an $\mathbf{NP}^{\mathbf{PosSLP}}$ algorithm. Thanks to [2], we obtain an upper bound of $\mathbf{coNP}^{\mathbf{PP}^{\mathbf{PP}^{\mathbf{PP}}}}$ for Positivity (see [17] for details). ◀